

Correction

Open Access

Digital PCR provides sensitive and absolute calibration for high throughput sequencing

Richard A White III[†], Paul C Blainey[†], H Christina Fan and Stephen R Quake^{*}

Address: Department of Bioengineering at Stanford University and Howard Hughes Medical Institute, Stanford, California 94305, USA

Email: Richard A White - raw937@sbcglobal.net; Paul C Blainey - blainey@stanford.edu; H Christina Fan - chfan@stanford.edu; Stephen R Quake^{*} - quake@stanford.edu

^{*} Corresponding author [†]Equal contributors

Published: 19 November 2009

Received: 11 August 2009

BMC Genomics 2009, **10**:541 doi:10.1186/1471-2164-10-541

Accepted: 19 November 2009

This article is available from: <http://www.biomedcentral.com/1471-2164/10/541>

© 2009 White et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Correction

After this article [1] appeared online, an error was called to our attention. The "universal probe" sequence UPL #149 in Table 6 appears with the 5' and 3' ends reversed. The correct sequence of this locked nucleic acid (LNA) probe is 5'-TCGCCGCC-3'. This typographical error does not affect any of the conclusions drawn in the article.

References

1. White RA, Blainey PC, Fan HC, Quake SR: **Digital PCR provides sensitive and absolute calibration for high throughput sequencing.** *BMC Genomics* 2009, **10**:116.