

RESEARCH ARTICLE

Open Access

# Analysis of microRNA transcriptome by deep sequencing of small RNA libraries of peripheral blood

Candida Vaz<sup>1</sup>, Hafiz M Ahmad<sup>2</sup>, Pratibha Sharma<sup>3</sup>, Rashi Gupta<sup>1</sup>, Lalit Kumar<sup>3</sup>, Ritu Kulshreshtha<sup>2</sup> and Alok Bhattacharya\*<sup>1,2</sup>

## Abstract

**Background:** MicroRNAs are a class of small non-coding RNAs that regulate mRNA expression at the post-transcriptional level and thereby many fundamental biological processes. A number of methods, such as multiplex polymerase chain reaction, microarrays have been developed for profiling levels of known miRNAs. These methods lack the ability to identify novel miRNAs and accurately determine expression at a range of concentrations. Deep or massively parallel sequencing methods are providing suitable platforms for genome wide transcriptome analysis and have the ability to identify novel transcripts.

**Results:** The results of analysis of small RNA sequences obtained by Solexa technology of normal peripheral blood mononuclear cells, tumor cell lines K562 and HL60 are presented. In general K562 cells displayed overall low level of miRNA population and also low levels of DICER. Some of the highly expressed miRNAs in the leukocytes include several members of the let-7 family, miR-21, 103, 185, 191 and 320a. Comparison of the miRNA profiles of normal versus K562 or HL60 cells revealed a specific set of differentially expressed molecules. Correlation of the miRNA with that of mRNA expression profiles, obtained by microarray, revealed a set of target genes showing inverse correlation with miRNA levels. Relative expression levels of individual miRNAs belonging to a cluster were found to be highly variable. Our computational pipeline also predicted a number of novel miRNAs. Some of the predictions were validated by Real-time RT-PCR and or RNase protection assay. Organization of some of the novel miRNAs in human genome suggests that these may also be part of existing clusters or form new clusters.

**Conclusions:** We conclude that about 904 miRNAs are expressed in human leukocytes. Out of these 370 are novel miRNAs. We have identified miRNAs that are differentially regulated in normal PBMC with respect to cancer cells, K562 and HL60. Our results suggest that post-transcriptional processes may play a significant role in regulating levels of miRNAs in tumor cells. The study also provides a customized automated computation pipeline for miRNA profiling and identification of novel miRNAs; even those that are missed out by other existing pipelines. The Computational Pipeline is available at the website: [http://mirna.jnu.ac.in/deep\\_sequencing/deep\\_sequencing.html](http://mirna.jnu.ac.in/deep_sequencing/deep_sequencing.html)

## Background

Small non-coding RNAs participate in a variety of processes from cell development and differentiation, stress responses to carcinogenesis by regulating gene expression [1-4]. Regulatory non-coding RNAs have been reported from almost all organisms from bacteria to mammals [5]. Among the various classes of non-coding

small RNAs (sRNAs), the most conserved and prominent ones are the microRNAs or miRNAs [for a recent review see, [6]]. Mature miRNA sequences are single stranded, typically 18-24 nucleotides long and encoded as a precursor molecule of about 60-120 nucleotides (in humans) [7]. These precursors are derived from processing pri-miRNA (usually in kilobases) by a ribonuclease, such as DROSHA [8]. Pre-miRNAs are also further cleaved to generate active mature miRNAs with the help of DICER [9]. So far more than 800 miRNAs have been described in

\* Correspondence: [alok0200@mail.jnu.ac.in](mailto:alok0200@mail.jnu.ac.in)

<sup>1</sup> School of Information Technology, Jawaharlal Nehru University, New Delhi, India

Full list of author information is available at the end of the article

human [10]. miRNAs interact with 3'UTR of mRNAs through base pairing and bring about their degradation, destabilization, or repression of translation through RISC, a complex of multiple proteins and miRNA-mRNA adduct [11,12]. Occasionally it can also up regulate gene expression [13].

The expression profiles of miRNAs have been determined in order to understand the role of miRNAs in a specific biological process [14]. The profiles generated revealed altered levels of miRNAs in different systems, such as oncogenesis and development [15,16]. These studies showed strong correlation between specific miRNA expression and phenotype suggesting it can be used as a potential biomarker for diagnosis and prognosis in human cancer [17]. Several groups of miRNAs have been identified that regulate the expression of tumor-associated genes while others seem to hold prognostic value in predicting patient survival. For example, miR-21 is frequently over expressed in various cancers [18]. It is now considered as an oncomiR that acts by down regulating PTEN, a tumor suppressor gene [19]. Similarly let-7 family of miRNAs is frequently down regulated in lung cancers [20]. miR-15a and 16-1 are often deleted in chronic lymphocytic leukemia [21]. These miRNAs target RAS, HMGA2 and BCL2 oncogenes, respectively, thereby regulating tumorigenesis [22-24]. Expression of let-7a, miR-210 and miR-200 clusters was demonstrated to be a strong prognostic marker for lung cancer, breast cancer and ovarian cancer, respectively [25-27]. Many of the fundamental processes are regulated by miRNAs, such as cell differentiation (miR-223, miR-145) [28,29], apoptosis (miR-34, miR-16) [21,30], body patterning (miR-9, miR-196) [31,32], nervous system and muscle development (miR-134, miR-1 and miR-133) [33,34].

A number of different approaches have been used for expression profiling, such as northern analysis [35], cloning [36], real time polymerase chain reaction [37], microarray analysis [38,39] and RNase protection assay [25,40]. These methods are not generally useful for discovery and expression profiling of low-abundance transcripts or yet unidentified novel miRNAs. Recently a number of different platforms have been developed for carrying out large scale parallel sequencing in order to generate genome wide sequences in a short time and reduced cost [41-43]. These have also been applied to analyze transcriptome including sRNA sequences [41]. Generally sequences derived using these platforms can be processed to generate expression profiles of known genes and suitable computational methods can be employed to identify unknown genes. Some of these platforms along with custom computation pipelines have been used to study sRNAs from different systems, such as plants, human embryonic cells and developing chicken embryo [44-46]. All these studies not only generated expression

profiles of known miRNAs but also identified a few novel miRNAs. Morin *et al.* used this strategy in combination with RNAfold, MiPred and an in house SVM model to identify novel miRNAs [47]. Though the study identified 104 novel miRNAs there was no attempt to verify these by other experiments. A software pipeline miRDeep has been developed to analyze large-scale sRNA sequencing data and identification of novel miRNAs [48]. Some of the filters used in this approach are highly stringent and it is likely that many novel miRNAs may be missed. Generation of expression profiles and identification of novel miRNAs from deep sequencing is dependent on tools used for analysis. It appears from this discussion that there is a need for development of efficient automated pipelines for analysis of deep sequencing data and setting up validation pipeline for checking the results, as newer algorithms may allow improved profiling and identification of novel miRNAs.


In this manuscript we have profiled sRNA expression from normal human peripheral blood mononuclear cells and two cancer cell lines using Solexa technology and developed automated computation pipelines for analyzing quantitative expression. Our pipelines use statistical analysis of the Solexa sequences for generation of expression profiles and a number of different methods for prediction of novel miRNAs. We believe that these pipelines are highly robust and can be useful for other studies.

## Results

### Sequencing and Annotation of Small RNAs

sRNAs were size selected by gel electrophoresis and then sequenced using a Solexa platform (Illumina, USA). The details of sequencing reads of the four samples are given in Additional file 1. The length of an average read was about 33-35 nucleotides. The processing of the raw sequences through computation pipeline is outlined in Figure 1 and described in "Methods". Briefly, the adaptor sequences were first removed from the sequence reads. Only those reads that were greater than 10 nucleotides were considered for further analysis. These sequences were clustered on the basis of sequence similarity and subjected to similarity searches using specific databases (rRNAs, tRNA, sn/snoRNAs, miRNAs, other non-coding RNAs). The reads that did not match known RNA sequences were checked to see if these were encoded by intergenic, intronic or exonic regions of the human genome (Figure 2).

Sequencing reads derived from mRNAs and rRNAs made up about 2-4 and 2-13% of the total reads of all the samples, respectively except that from K562 cells. Since mRNAs and rRNAs are likely to have been derived by degradation it appears that the degradation was minimal in these samples. The sequencing reads derived from miRNAs were estimated to be about 60-80%, similar to





some of the other studies on sRNAs [47]. The result from K562 was quite different. Only about 18% of total sRNA population was found to be derived from miRNA. Low

level of miRNAs in K562 cells is unlikely to be due to artifact as we got similar values in two independent experiments (data not shown). Out of a total of 904 miRNAs (718 major and 186 star sequences) present in miRBase version 14, 534 miRNAs were detected in at least one of the four samples sequenced [49]. A total of 370 miRNAs were not detected in any of the samples.

### MicroRNA expression patterns

Absolute sequence reads were transformed into transcript abundance by first normalizing the data in 'transcripts per million (TPM)' for each library (see "Methods"). The expression levels ranged from less than 10 to more than 100,000 counts (Figure 3, refer Additional file 1). Thus the sequencing data revealed a wide range of expression levels spanning five orders of magnitude. Several members of the let-7 family, 103, 185 and 320a were observed as some of the highly expressed miRNAs (Figure 4). These miRNAs were also reported as highly abundant miRNAs in peripheral blood mononuclear cells using "Taqman microRNA assay" [50]. While the members of let-7 family, let-7a, f and g represented about 77% of the total miRNA counts, a number of miRNAs with less than 10 counts including singletons were also noticed (Additional file 1). It is not clear if the singletons represent true transcript or noise of the system. miR-219-5p was one of the singletons found in all the four samples suggesting that such sequences may not be due to experimental noise or a chance event. Star sequences of many miRNAs (20% of all known miRNAs) were also observed mostly at low abundance level. Some of the highly abundant star sequences observed by us were miR-106-b\* (N1, TPM- 288.28); miR-17\* (N1, TPM- 1321.41); miR-92a-1\* (N1, TPM- 942.78), miR- 25\* (N1,





TPM-265.44), and miR-374a\* (N1, TPM-201.34). The results suggest that sRNA sequencing is a good approach for studying miRNA\* abundance.

### Expression patterns of miRNA clusters


miRNAs are often present in gene clusters and it is generally believed that individual miRNAs are generated from common polycistronic transcript by post-transcriptional processing [51,52]. Since the genes in clusters are co-regulated and co-transcribed it is expected that the levels of these miRNAs are likely to be similar. This was investigated by checking the relative levels of individual miRNAs present in clusters using the expression patterns derived from deep sequencing. A total of 265 miRNA genes organized in 68 clusters were studied. As the sequencing reads represent mainly mature miRNAs and not individual precursors, it was not possible to analyze some of the clusters, such as those containing miR-15a, 16-1 and 15b, 16-2 that encode different precursors but


the same mature miRNA. Furthermore, 19 clusters were also excluded from analysis as the miRNAs encoded in these clusters were either not expressed or expressed below 1 TPM based on our data. Therefore in this study results from analysis of only 20 clusters are shown (Additional file 2).

The level of expression of different miRNAs in many clusters displayed variable expression as exemplified by the cluster containing miR-532 and 99b (Figure 5). In some cases the variation observed within a cluster was as much as 500 fold and the miRNA that showed highest level of expression was not the one closest to the transcription start site. For example, miR-25 showed highest expression though it was farthest from the start site in miR-106b cluster (Figure 5). This is likely to be due to variation in processing mechanisms at post-transcriptional level as has been seen in case of p53-mediated alteration in processing some of the miRNAs, such as miR-16-1 [53]. Overall the results suggest that the levels of miRNAs in any cell may be regulated by a number of different processes and mechanistic details of many of these are not yet known.


### Differentially Regulated miRNAs in K562 and HL60

Microarray based methods have been used extensively to profile expression levels of annotated genes at genomic scale. These methods cannot be used satisfactorily for analyzing expression levels of unknown transcripts. Deep sequencing of cellular RNAs is an alternative approach for deciphering expression profiles of genes [46,54]. Though a few reports describing analysis of differentially expressed miRNAs deciphered by deep sequencing are available, there is no study yet on human PBMC. Moreover, there is no standardized tool available for profiling miRNA expression using deep sequencing of sRNA [47,55]. Therefore we have studied normal peripheral blood mononuclear cells from two different individuals and cancer cells of myeloid lineage, K562 (chronic myelocytic leukemia) and HL60 (acute promyelocytic leukemia) by deep sequencing of sRNAs followed by analysis of the sequences using a custom designed computation pipeline. We used intersection of two methods for finding differentially expressed genes; a) SAM (significance analysis of microarrays) that allows for the control of false detection rate (FDR) and b) fold change, that is, miRNAs that showed more than 2.5 fold differences when compared to both normal samples. A list of differentially regulated miRNAs is shown (Figure 6). Interestingly, miR-1, 101, 106b, 146b-5p, 151-3p, 192, 21, 22, 27b, 30e and 361-3p displayed low expression in both the cell lines indicating a common role in leukemia genesis or progression (Figure 6, see inset). In K562 most of the miRNAs were down regulated except miR-486-3p and miR-504. The expression patterns derived from sRNA sequences were





**Figure 5** Differential expression of individual miRNAs present in the same cluster in different datasets. Here TPM (transcript per million) is used as a measure of expression. [A] miRNAs belonging to cluster miR-532, [B] cluster miR-99b and [C] cluster miR-106b in normal PBMC, K562 and HL60 cell lines. A large variation in expression levels of different miRNAs present within the same cluster is observed.




validated by RNase protection assay (RPA) which allowed both qualitative and quantitative analysis of RNA levels [25,56]. Down regulation of expression of some of the miRNAs, such as miR-16, 22, 27a, 192 and let-7g in CML cell line K562 was observed using RPA assay (Figure 7). RNU6B was used as a loading control, for all the samples. We have also used quantitative RT-PCR for validating some of the results derived by deep sequencing and RPA. Fold differences in the expression of some of the differentially expressed genes in normal and K562 cells (miR-22 and miR-27a) using all three methods were found to be comparable indicating quantitative nature of these approaches (Additional file 3).


While miR-21 was found to be one of the down regulated miRNAs in our study, it is expressed at high level in all the cancers tested till date indicating that miRNA expression is likely to be cell type and context dependent

[18]. Reduced expression of some of the miRNAs, such as miR-16, 151 and 142 was previously reported in CML [57-59]. Up regulation of the polycistronic miR-17-92 cluster in CD34+ cells of CML patients was initially described by Venturini *et al.*, 2007 and later was challenged by Agirre *et al.*, 2008 [59,60]. According to the latter group there is no significant increase in the expression of this cluster in CML cells. We too did not observe any induction of this cluster in K562 cell line in agreement with the latter group.

A significant number of differentially expressed miRNAs of HL60 (downregulated-miR-101, 126, 27b, 7; up regulated- let-7a, let-7d, miR-181a, -181a\*, -181b and miR-199b) were mapped to chromosome 9 (Figure 8). Differential expression of miR-181a family (miR-181a, 181a\* and 181b) was previously reported [61]. Some of the differentially regulated miRNAs of K562, such as



miR-101, miR-27b and miR-24 were also encoded by chromosome 9. The genomic locations of some of these miRNAs were distant from the ABL gene locus. Alterations in chromosome 9 are associated with a large number of diseases, particularly cancer [62]. Therefore, there



may be a link between alterations in chromosome 9 and differential expression of miRNAs.

#### Predicted targets of differentially regulated miRNAs

Identification of putative targets may help to understand the biological role of the differentially regulated miRNAs. In this study two different approaches were used for target identification. In the first approach a number of different software tools were employed for target prediction and in the second approach mRNA expression profiles were generated and the genes that showed inverse correlation with miRNA expression were identified. The second approach is based on a number of observations where target mRNA degradation by miRNAs were reported [63,64]. For computational prediction stringent criteria, that is, only those genes that are predicted to be target of specific miRNAs by at least five of the eleven established target prediction programs (DIANA-microT, MicroInspector, miRanda, MirTarget2, miTarget, NBmiRTar, PicTar, PITA, RNA22, RNAhybrid, and TargetScan/TargetScanS) integrated by miRecords and also showed expression levels that are inversely correlated with the specific miRNA levels were considered [65]. Putative targets of differentially regulated miRNAs are shown in Additional file 4 (K562) and Additional file 5 (HL60). Genes targeted by three or more miRNAs are shown in Table 1. Genes that are targets of multiple miRNAs are likely to be tightly regulated and may show graded response on the basis of expression of different miRNAs [66]. MEIS2, up regulated in K562, was found to be the target of up to five different miRNAs, down regulated in both K562 and HL60. MEIS2 has been previously reported to be important for myeloid leukemogenesis [67]. Similarly, SMAD7 a down regulated gene in HL60 is predicted to be a target of three miRNAs, up regulated in HL60. It is known that SMAD7 regulates SMAD and mitogen activated kinase (MAPKs) signaling and controls erythroid and megakaryocytic differentiation of erythroleukemia cells [68]. TRIB2, multiply targeted by three up regulated miRNAs in HL60 has been reported to be present at low levels in AML [69].

#### Expression of intronic miRNAs

Many intron-encoded miRNAs are processed in DROSHA independent way (miRtrons) [70]. We have also analysed expression levels of differentially regulated intronic miRNAs. In order to check if differential expression of intronic miRNAs is due to transcriptional or post-transcriptional mechanisms, the levels of intronic miRNAs were compared with that of the host transcripts. Comparative analysis revealed positive correlations with respect to 8 out of 14 intronic miRNA - mRNA pairs (Table 2). Unlike miRNAs in clusters, the expression of

**Table 1: List of Genes targeted by 3 or more differentially regulated miRNAs in K562 and HL60.**

Genes targeted by 3 or more downregulated miRNAs in K562		Genes targeted by 3 or more downregulated miRNAs in HL60		Genes targeted by 3 or more upregulated miRNAs in HL60	
<b>ABCC5</b>	let-7g, let-7i, miR-101	<b>BCL11A</b>	miR-1, miR-21, miR-152	<b>ARID3B</b>	let-7a, let-7c, let-7d, miR-98
<b>ACVR2B</b>	miR-16, miR-101, miR-186,	<b>EDEM3</b>	miR-27b, miR-30e, miR-101	<b>ARRDC3</b>	miR-25, miR-181a, miR-181b
<b>ADAMTS3</b>	miR-16, miR-30e, miR-146b-5p	<b>GALNT7</b>	miR-7, miR-27b, miR-30e	<b>BAZ2A</b>	miR-25, miR-99a, miR-181a
<b>ARRDC4</b>	miR-27a, miR-27b, let-7i, let-7g	<b>MEIS2</b>	miR-27b, miR-30e, miR-101, miR-194	<b>CCR7</b>	let-7a, let-7b, let-7c, let-7d
<b>BZW2</b>	let-7g, let-7i, miR-101	<b>PHF6</b>	miR-1, miR-30e, miR-101, miR-106b	<b>CD200R1</b>	let-7a, let-7b, let-7c, let-7d, miR-98
<b>CDC25A</b>	let-7g, let-7i, miR-16, miR-21, miR-339-3p	<b>PTGFRN</b>	miR-30e, miR-106b, miR-146b-5p	<b>CPEB2</b>	let-7a, let-7b, let-7c, let-7d, miR-25, miR-98
<b>CIT</b>	miR-27a, miR-27b, miR-30e, miR-142-5p	<b>SATB2</b>	miR-22, miR-27b, miR-30e	<b>CPEB4</b>	miR-25, miR-181a, miR-181b
<b>CLCN3</b>	miR-1, miR-101, miR-27a, miR-27b			<b>DUSP16</b>	let-7a, let-7b, let-7c, miR-98
<b>CLDN12</b>	let-7g, let-7i, miR-16			<b>EGR3</b>	let-7a, let-7b, let-7c, let-7d, miR-98, miR-181a
<b>CPEB1</b>	let-7g, let-7i, miR-1, miR-22			<b>EPHA4</b>	let-7a, let-7b, let-7c, miR-98, miR-181a
<b>DCBLD2</b>	miR-16, miR-24, miR-101			<b>FOS</b>	miR-181a, miR-181b, miR-221
<b>E2F7</b>	miR-16, miR-27a, miR-27b			<b>GCNT4</b>	let-7a, let-7b, let-7c, let-7d, miR-98
<b>GALNT7</b>	miR-27a, miR-27b, miR-30e			<b>GPX7</b>	let-7a, let-7b, let-7c
<b>HIC2</b>	let-7g, let-7i, miR-24, miR-30e, miR-146b-5p			<b>IGF2BP2</b>	let-7a, let-7b, let-7c, miR-98, miR-181a
<b>IGF2BP3</b>	let-7g, let-7i, miR-142-5p			<b>ITGB3</b>	let-7a, let-7b, let-7c, miR-98
<b>LIN28B</b>	let-7g, let-7i, miR-30e			<b>KIAA1539</b>	let-7a, let-7b, let-7c, let-7d, miR-98
<b>MEIS2</b>	let-7i, miR-27a, miR-27b, miR-30e, miR-101			<b>KLHL24</b>	let-7a, let-7b, let-7c, miR-98, miR-618
<b>NEDD4</b>	miR-27a, miR-27b, miR-30e			<b>LRIG1</b>	let-7a, let-7b, let-7c, let-7d, miR-98, miR-425
<b>PDIK1L</b>	miR-1, miR-16, miR-22, miR-142-5p			<b>MAP3K3</b>	let-7b, miR-181a, miR-181b
<b>PHF6</b>	miR-1, miR-30e, miR-106b, miR-186			<b>MYO1F</b>	let-7a, let-7b, let-7c, let-7d
<b>PTGFRN</b>	miR-30e, miR-106, miR-146b-5p			<b>PPP1R16B</b>	let-7a, let-7b, let-7c, let-7d, miR-98
<b>RELN</b>	miR-16, miR-27a, miR-27b			<b>PRKCE</b>	miR-25, miR-181a, miR-181b
<b>SATB2</b>	miR-16, miR-22, miR-27a, miR-27b, miR-30e			<b>RAB11FIP4</b>	let-7a, let-7b, let-7c, let-7d, miR-98
<b>SGMS1</b>	miR-27a, miR-27b, miR-106b, miR-142-5p			<b>SLAMF6</b>	let-7a, let-7b, let-7c, let-7d, miR-98
<b>SPRED1</b>	miR-1, miR-16, miR-101			<b>SLC30A4</b>	let-7a, let-7b, let-7c, miR-98
<b>STRBP</b>	let-7g, let-7i, miR-146b-5p			<b>SLC35D2</b>	let-7a, let-7b, let-7c, let-7d, miR-98
<b>WEE1</b>	miR-16, miR-27a, miR-106b			<b>SLC4A4</b>	let-7a, let-7b, let-7c, let-7d
				<b>SMAD7</b>	miR-25, miR-181a, miR-181b
				<b>SNN</b>	let-7a, let-7b, let-7c, let-7d, miR-25, miR-98, miR-181a
				<b>STK40</b>	let-7a, let-7b, let-7c, let-7d, miR-98
				<b>SYT11</b>	let-7a, let-7b, let-7c, let-7d
				<b>TMEM2</b>	let-7a, let-7b, let-7c, let-7d
				<b>TRIB2</b>	let-7a, let-7b, let-7c, miR-98
				<b>UTRN</b>	let-7a, let-7b, let-7c, let-7d, miR-98

**Table 2: Correlation of expression patterns in A) K562 and B) HL60 cancer lines between differentially regulated intronic miRNAs and their host genes.**

A) K562		
miRNA	Host Transcript	miRNA:Host transcript Status
miR-342	EVL	Downregulated
miR-548e	SHOC2	Downregulated
miR-486	ANK1	Upregulated

B) HL60		
miRNA	Host Transcript	miRNA:Host transcript Status
miR-22	C17orf91	Downregulated
miR-151	PTK2	Downregulated
miR-199b	DNM1	Upregulated
miR-25	MCM7	Upregulated
miR-618	LIN7A	Upregulated

some of the intronic miRNAs may be transcriptionally controlled.

#### miRNA Biogenesis Machinery

The analysis of microarray data revealed induction in the expression of some of the miRNA biogenesis genes (RNASEN, DGCR8, XPO5, RAN) in K562 cell line (Table 3). However, DICER1 was found to be down regulated (1.79 fold). This would result in a likely increase in the pre-miRNA population in cytosol but decrease in accumulation of mature miRNAs. This is consistent with the observation that K562 cells contain relatively less amount of miRNAs and that 23 out of 25 differentially expressed miRNAs are down regulated (see Figure 6). This observation was not due to artifact as two independent sequencing of sRNAs from K562 cells gave similar results (data not shown here). Low DICER1 and let-7 levels were also observed in lung adenocarcinoma [20]. In contrast, a higher level of DICER1 in prostate adenocarcinoma accounted for up regulation of 39 of 45 differentially expressed miRNAs [71]. In HL60 some of the compo-


nents of biogenesis machinery (RNASEN and XPO5) were found to be up regulated.

#### Identification of Novel MicroRNA genes


In principle deep sequencing of sRNAs should generate sequences from as yet unannotated regions of the genome. The analysis of the sequences through computational pipeline (Figure 9) showed a large number of unannotated sequences that are encoded by either intergenic or intronic regions. Since miRNAs are predominantly encoded by intergenic and intronic regions, these sequences were analyzed by a set of computational tools to identify putative novel miRNAs. Since predictions are based on identifying miRNA precursors, genomic regions (70 nucleotides) surrounding the sRNA sequences were extracted (see "Methods" for details). A total of 370 (357 major + 13 minor) novel miRNAs were predicted using computational pipeline (Additional file 6). The sequences and chromosomal locations of the predicted novel miRNAs from intronic and intergenic regions are listed in the supplementary table (Additional files 7 and 8). More than

**Table 3: Altered levels of miRNA biogenesis and miRISC components in K562 and HL60.**

Protein	Function	K562/Normal (Fold Change)	HL60/Normal (Fold Change)
DROSHA (RNASEN)	PrimiRNA processing	2.238954	2.017825
DGCR8	PrimiRNA processing	2.078018	Unchanged
XPO5	Exporting premiRNA	2.669646	2.199101
RAN	Exporting premiRNA	1.832214	Unchanged
DICER1	PremiRNA processing	-1.79971	Unchanged



95% of the novel miRNAs showed frequency counts lower than 10. Many novel miRNAs in our dataset showed presence of 1-11 isomiRs with varying frequencies. For example, frequencies ranging from 1-280 were observed for eight isomiRs of jnuhsa-204. Presence of isomiRs certainly strengthens classification of these molecules as novel miRNAs [47]. These datasets were also analyzed using miRDeep, a published tool for miRNA analysis of deep sequencing data [48]. The number of novel miRNAs predicted by miRDeep is much less than that using the pipeline described here (Additional files 7 and 8). This may be due to stringent criteria used by miRDeep that quite often misses known miRNAs. The expression of a few novel miRNAs, missed by miRDeep but predicted by "in-house tools", was validated through real-time PCR (for precursor miRNA) or RNase protection assay (for mature miRNAs) as shown in Figure 10 and 11. Interestingly miRNAs that were present as singleton (for example- jnuhsa-4-3p, jnuhsa-93-3p) were also detected using RNase protection assay (Figure 11). We also mapped predicted novel miRNAs to clusters based on the inter-miRNA distance of less than 20 kb (Figure 12). Some of the clusters contained only novel miRNAs while many others were part of previously known clusters




[72]. Based on miRDeep prediction a total of 12 novel miRNAs showed similarity in the seed region with the known miRNAs indicating that these miRNAs are likely to belong to the same family and thus may share common biology (Additional files 7 and 8).

## Discussion

Our effort to understand the microRNAome of human leukocytes through deep sequencing technology has given us several interesting observations. These include discovery of novel miRNAs and a valuable list of differentially expressed miRNAs in chronic and acute myelogenous leukemia. The diversity in the miRNA wealth was realized through the detection of large number of novel miRNAs. Our major findings are a list of expressed miRNAs (534 known and 370 novel) in leukocytes and the discovery that the expression of miRNAs may be controlled by regulating post-transcriptional events, such as manipulating the level of DICER, an enzyme involved in biosynthesis of miRNAs in tumor cells (K562). Generally, singletons in the sequenced samples are not taken into consideration, as these are likely to occur due to potential sequencing error or transcription noise [73]. Our experimental results show convincingly that such miRNAs are likely to be real and not a result of artifact of sequencing.

The known miRNAs found to be most abundant in all the four samples were members of the let-7 family, 103, 185 and 320a, which were also reported in peripheral blood mononuclear cells using "Taqman microRNA assay" [50]. Therefore our results are in agreement with previous studies based on a different methodology. There are multiple mechanisms that are likely to regulate miRNA levels similar to that of mRNAs. These include





isomiRs affect the target recognition needs to be evaluated.

A number of differentially expressed miRNAs and their potential targets were identified by comparing miRNA and mRNA levels of normal PBMC versus that of two myeloid cancer cell lines K562 and HL60 along with computation approach. This strategy was used as degradation of many mRNAs was observed as a result of miRNA action [77]. Both K562 and HL60 have been used as reference cell lines for studying the cellular and molecular events involved in the proliferation and differentiation of chronic myeloid or acute promyelocytic leukemia, respectively. Many miRNAs that were previously reported to be involved as tumor suppressors, such as let-7 g, miR-101, 16 and 192 were found to be down regulated in the leukemia cell lines. Many of the differentially regulated miRNAs, identified in this study have been reported to be involved in various cellular processes like cell cycle (miR-192) [78], apoptosis (miR-16, -126, -98) [21,79,80] differentiation (miR-27a, -181, -342, -223) [28,81-83] DNA repair (miR-24, -210) [84,85] metastasis [86], erythroid maturation (miR-22) [87], erythropoiesis (miR-24) [88] and hematopoiesis (miR-142, -181) [89,90]; highlighting their putative role in leukemogenesis or progression. Strikingly, majority of the differentially regu-

lated miRNAs were found to be down regulated in K562 cell line. It is possible that low level of DICER, as revealed by gene expression profiling, is responsible for an overall reduction in the miRNA population. Global down regulation of miRNAs in mouse T cells using PCR and conventional sequencing approach was reported before [91]. However, the cause of the down regulation was not clear, as mRNA profiling was not carried out. DICER has also been implicated in mouse T cell functions [92]. Our studies strongly suggest that there may be a link between DICER and miRNA levels. Many of the genes, identified in this study as the potential targets of differentially regulated miRNAs are known to be involved in cancer through their effects on cell differentiation (CDK6, LIFR), apoptosis (PIM1) or hematopoiesis (GATA2, TAL1) [93-97]. Interestingly, majority of the targets of the cancer-associated miRNAs, such as -ARHGEE, CDK6, ETV5, GATA2, PIM1, LIFR, TAL1, PPARG, RANBP17, TFRC FOXP1 LPP, NCOA2 and NR4A3 have been previously shown to be associated with leukemogenesis.

One of the features of this study is the computation pipeline that identifies novel miRNAs among sRNA sequences. This pipeline is different from other available pipelines as it has custom designed tools for identification of novel miRNAs that involve pre-processing of the

sequences, an exhaustive elimination pipeline, folding and filtering to get novel predictions. The evidences supporting the predictions are identification of potential precursors by specific precursor identification tools, presence of star sequences, presence of isomiRs and in some cases experimental validation. The fact that even a single sequence read may not be a noise or error, has been validated by observing corresponding sRNA experimentally and indicates the power of our computation pipeline in identification of novel miRNAs. The total number of novel miRNAs identified from all the four samples is 370. Therefore, total number of miRNAs encoded by the human genome may be much larger than current estimate [98].

## Conclusions

In conclusion we list about 904 miRNAs that are expressed in normal and cancerous leukocytes, nearly 41% of these are novel. Our analysis shows that some of the novel miRNAs are likely to be clustered in the genome similar to many known clusters of miRNAs. Moreover, we have also identified a number of miRNAs that are differentially expressed in cancer cells studied by us. These are likely to be new markers/cause of carcinogenesis. In addition we show that miRNA levels can be regulated at post-transcriptional processing stage. Our results also show that singletons in deep sequencing reads are unlikely to be sequencing artifacts. In K562 cells low levels of miRNAs is correlated with reduced level of the enzyme DICER. In addition we have developed an improved automated computation pipeline for analysis of deep sequencing data to obtain quantitative profiles of known and novel miRNAs. Based on our observations, it is now possible to generate leukocyte specific miRNA arrays in order to study expression profiles of all miRNAs relevant to leukocytes. Similar approach can be used to generate other tissue specific miRNAome.

## Methods

### Cell lines and blood samples

The human chronic myeloid leukemia blast crisis cell line, K562 and acute promyelocytic leukemia cell line, HL60 were obtained from National Centre for Cell Sciences, Pune and maintained in RPMI 1640 and DMEM (Gibco) medium, respectively. The medium was supplemented with 10% FBS and penicillin-streptomycin and maintained at 37°C with 5% CO<sub>2</sub> in incubator chamber. Buffy coat of healthy blood donors (N1 and N2) were collected from volunteers. Red cell lysis buffer (0.144M NH<sub>4</sub>Cl, 0.01M NH<sub>4</sub>HCO<sub>3</sub>) was added to buffy coat to lyse the remnant RBCs and pure WBC population was obtained by centrifugation at 3000 g.

### RNA Isolation and sequencing

Total RNA isolation was carried out from peripheral blood and cell lines using TRIzol<sup>®</sup> Reagent (Invitrogen) as per manufacturer's instruction. RNA preparations were stored at -80°C till further use. Small RNA (sRNA) population was isolated by separating 10 µg of total RNA on denaturing polyacrylamide gel electrophoresis (PAGE) and cutting a portion of the gel corresponding to the size 18-30 nucleotides based standard oligonucleotide markers. Adapter (5') was ligated to sRNA population and ligated RNAs (40-60 nt) were purified by running on urea PAGE. This was followed by 3' adapter ligation and purification of adapter ligated RNAs (70-90 nt) in a similar manner. Modified sRNAs were reverse transcribed and then PCR amplified with adapter specific primers and the amplified cDNAs were finally purified on Urea PAGE to generate cDNA tag libraries for sequencing by illumina genome analyzer. The average number of sequencing reads was around 4.9 million.

### Datasets

Four sRNA sequencing data comprising of peripheral blood leukocytes of two normal individuals (N1, N2) and tumor cells K562 and HL60 were obtained from illumina fast track sequencing services. For each sample a Sequence file and a Tag file was provided. The Tag file comprises of unique sequences with their corresponding frequency. Tag Files are generated post alignment as a summary of Sequence Files and every Sequence File has a corresponding Tag File. Tag Files are generated to give the researcher an indication of most common to most rare sequences in the dataset. The numerical frequency of each sequence in the Sequence File for gene expression gives a true indication of relative expression of sequence transcripts. The unique sequences in the Tag files contain a 3' adaptor sequence (TCGTATGCCGTCTTCTGCTTG). The amount of the 3' adaptor is variable and is dependent on the length of the sRNA. Each delivered sequence is 33/35 bases in length. A part of the adaptor sequence is seen in each sequence if the sRNA is shorter than 33/35 bases. This adaptor and segments of it needs to be trimmed for proper alignment to the transcriptome/genome.

### Preparation/Processing of the datasets

*i). Removal of the adaptor sequences:* Since the sequence of the adaptor is known, a perl script was written to trim the adaptors.

*ii). Clustering and removal of redundancy after removal of the adaptor sequence:* Although the tag file contained unique sequences, there were some, which after the removal of the adaptor were redundant. These identical sequences were represented once and their frequency was summed up.

**iii). Sequences less than 10 nucleotides in length were excluded**

**iv). Conversion into fasta format:** The final trimmed file was then converted into fasta format where the unique header (sequence identity) retained the information of the sequences length and frequency. The sequence ID comprised of a running number along with the length and frequency of that sequence.

#### Small RNA annotation

The sRNA sequences obtained were annotated against the known databases using the following protocol-

##### **i). Known/annotated sequence databases**

List of the databases used for the annotation/elimination pipeline were:

- a) **Mature miRNAs:** from miRNA registry, release 14 (includes a total of 904 miRNAs [718 mature (major) +186 stars (minors)]).
- b) **ncRNAs:** from Ensembl "Homo\_sapiens.NCBI36.56.ncrna.fa" (includes the precursor miRNAs and other ncRNAs like sn/sno/sca RNAs, tRNAs, rRNAs).
- c) **RNA database** from the FTP site NCBI (includes rRNAs and mRNAs).
- d) **The exons** were obtained from the Contig files by a self written script.
- e) **Intergenic/intronic sequences:** obtained through in-house built Perl script (using the *Homo Sapiens* Contig file 29 Feb, 2008 version). These sequences served as a source for finding novel miRNAs (intergenic/intronic).

##### **ii). Finding Known miRNAs**

One of our objectives was to study the expression pattern of the known miRNAs. To generate the expression profile of the known miRNAs, the sRNA sequences of all the 4 samples were matched against the known miRNA sequences using BLASTN and an in house built pattern matching tool. The parameters used for BLAST were tuned to obtain maximum matches, such as the word size was set to 7 nucleotides, filtering was turned off and the number of alignments reported were increased.

##### **iii). The Elimination Pipeline**

An in house - built script was written to do a fast matching of the sequences with the created databases. A mismatch of up to 2 nucleotides was allowed. The pool of unmatched sequences at the end of the pipeline served as a source of novel miRNAs (Figure 1).

#### Normalisation of the data/Calculation of Transcripts Parts Per Million (TPM)

Normalisation was carried out as the total number of reads from different experiments was not same and variations in the number of reads of individual miRNA can be due to sequencing depth. The number of reads of a tran-

script/sequence (representing a known miRNA) was divided by the total clone count of the sample and multiplied by  $10^6$ . The total clone count is the sum of the frequencies of all the unique sequences/transcripts present in the trimmed file. The Additional file 1 contains a list of the known miRNAs found in all the 4 samples along with their respective frequencies and TPM values.

#### Selecting Differentially expressed MicroRNAs

Differentially expressed miRNAs were identified by using a combination of two methods:

##### **i) SAM Analysis**

We performed the t-test procedure within significance analysis of microarrays (SAM) to look for differentially expressed miRNAs. SAM calculates a score for each gene on the basis of the change in expression relative to the standard deviation of all measurements.

##### **ii) Fold Change**

The known miRNAs in normal samples were compared to the cancer cell line. The miRNAs showing more than 2.5 fold difference as compared to both the normal cells were considered as differentially regulated. The samples with less than 10 TPM in both normal and cancer samples were ignored.

#### Novel miRNA Prediction

The strategy is based on first removing all known RNAs including those derived from exonic regions and then identifying those that are derived from intronic and intergenic regions. These were then subjected to some of the *ab initio* miRNA prediction algorithms like SCFG based CID-miRNA [99] and CSHMM [100] that use stringent criteria to distinguish between real and pseudo miRNAs. Besides this we ensure high discriminative power by using the following filters:

- a) Occurrence of the sequence in the stem region of the precursor
- b) Presence of IsomiRs
- c) Presence of minor/star sequences.
- d) Taking a consensus among the prediction tools

Following is the detailed pipeline used for the prediction of novel miRNAs:

##### **i) Extraction of matches from the intergenic/intronic regions of the human genome**

The unmatched sequences (from the elimination pipeline) were matched to the intergenic/intronic regions. The exact matched sequences were extracted along with 70 nucleotides flanking both the ends representing potential precursor sequences.

##### **ii). Folding the extended sequences and checking its location in the folded structure**

The sequences were scanned for presence of potential precursor miRNA using CID-miRNA [99] and CSHMM [100] prediction tools. The folded sequences generated

were then checked to see if the sRNA (putative mature miRNA obtained by sequencing) occurs in the folded putative precursor as the window scanning approach used could report a folded structure not involving the concerned sRNA. Only those hairpins were kept which contained sRNA. The next step involved locating the position of the sRNA in the hairpins. Since mature miRNAs are known to be arising from the stem portion and not the loop, only those hairpins in which the sRNAs occurred in the stem were classified as correct cases and the remaining as prediction errors. These correct cases were further tested by MiPred [101].

### **iii). Finding IsomiRs and Star sequences**

A list of all the predicted correct precursor sequences was created and the sRNAs derived from common precursors were grouped into a common family. The sRNAs derived from the same precursors were kept together in a family. The most abundant member was designated as the mature miRNA. The sRNAs that differed from the representative by a few nucleotides were called IsomiRs and those that had a different, partially complementary sequence and were located in the other strand (stem of the hairpin loop) were called stars. The Additional files 7 and 8 comprise of the novel miRNAs grouped into families on the basis of sRNAs falling within the same precursor. The representative sequence was chosen on the basis of abundance. The most abundant sequence was selected as representative of the family. The scores from the 4 tools (CID-miRNA, CSHMM, miRDeep, MiPred assigned to the corresponding precursors are also listed).

### **iv). Locating the potential - novel miRNA in the genome**

On the basis of the positions of the predicted novel miRNAs, their tendency to occur in a known cluster or a new cluster was checked. miRNAs located within 20 kb of the known or novel miRNA were considered as part of the same cluster.

### **v). Removal of the redundant miRNAs to get the final list of novel miRNAs**

The novel miRNAs from the 4 samples (both intergenic and intronic) were pooled and the redundancy was removed to get a final set of novel miRNAs. These were given a unique name (Additional file 6). The representative miRNAs in the Additional files 7 and 8 along with their sample IDs also have these names.

### **RNAse Protection Assay (RPA) for detection of mature miRNA and real time RT-PCR for precursor miRNA detection**

The RPA assay for mature miRNA detection was done using *mirVana*<sup>™</sup> miRNA detection kit as per supplier's instructions (Applied Biosystems). To detect the precursor miRNA, total RNA was treated with DNAase I (MBI fermentas). Reverse transcription was done using superscript<sup>®</sup> III (Invitrogen) reverse transcriptase as per supplier's instructions. A list of primers used for real time

RT-PCR amplification of precursors of novel miRNAs is given in Additional file 9. Real-time PCR was done with following parameters- Initial denaturation - 94°C for 2 min, denaturation- 94°C for 30 sec, annealing- 60°C for 1 min for 40 cycles using SYBR Green PCR Master Mix (Applied biosystems).

### **Microarray analysis**

Total RNA from normal PBMC and two myeloid leukemia cancer samples (K562 and HL60) was sent for gene expression profiling using Ocimum microarray platform (Ocimum, Hyderabad, India). The expression data for each sample was generated on Affymetrix Human Gene 1.0 ST arrays. A fold change cut-off of  $\pm 1.5$  resulted into FDR < 0.001 for both the comparisons. The quality control analysis was carried out using Affymetrix Expression Console (EC). The statistical analysis was performed using R-programming language and the biological analysis was carried out using Genowiz<sup>™</sup> software. The data obtained was normalized and genes showing more than 1.5 fold differences in the cancer cell lines as compared to normal were marked as differentially expressed genes. A total of 1856 genes were found to be up regulated and 1696 were down regulated in K562. In HL60, 1497 genes were up regulated and 1213 genes were found as down-regulated (Additional file 10).

### **miRNA target prediction**

The most probable targets of the differentially regulated miRNAs were fished out using following two criteria - 1. Prediction by at least five of the established target prediction programs- A list of putative targets was prepared (List A) using intersection between at least five of the eleven established target prediction programs compiled by miRecords: (DIANA-microT, MicroInspector, miRanda, MirTarget2, miTarget, NBmiRTar, PicTar, PITA, RNA22, RNAhybrid, and TargetScan/TargetScanS). 2. Inverse correlation in expression pattern between miRNA and coding genes- The putative target genes in list A was compared to the list of differentially regulated genes (showing more than 1.5 fold difference) (List B) in K562 and HL60 as obtained from expression profiling data, and only those genes that show inverse correlation to the miRNA levels were considered as most genuine putative targets of the select miRNAs.

### **Availability and Requirements**

The entire computational pipeline described in this paper is available at the website: [http://mirna.jnu.ac.in/deep\\_sequencing/deep\\_sequencing.html](http://mirna.jnu.ac.in/deep_sequencing/deep_sequencing.html). The software is also available on request.

### **Accession numbers**

The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus and are accessi-

ble through GEO Series accession number GSE19833 Super series: <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE19833>

The GSE19833 Super series comprises of the following sub series:

Microarray Expression data: GSE 19789 (consists of raw data and normalized data) Deep sequencing data: GSE 19812 sequence data (consists of the raw sequences, untrimmed tag and trimmed tag files and the expression profile of the known miRNAs (Additional File 1).

## Additional material

**Additional file 1 The known miRNAs expression pattern for all the 4 samples.** The known miRNAs found in the 4 samples with their corresponding frequency and TPM values.

**Additional file 2 miRNA cluster analysis.** Cluster analysis of the known miRNAs in Normal1, Normal2, K562 and HL60.

**Additional file 3 Comparison of the sensitivity of the three miRNA detection methods (Deep sequencing reads, RNase Protection assay and quantitative RT-PCR).** Fold change differences in normal PBMC versus K562 is presented for miR-27a and miR-22 using the three transcript detection methods.

**Additional file 4 List of predicted targets of differentially regulated miRNAs showing inverse correlation in microarray data in K562.**

**Additional file 5 List of predicted targets of differentially regulated miRNAs showing inverse correlation in microarray data in HL60.**

**Additional file 6 List of the predicted 370 novel miRNAs.** The precursor and mature sequences of the total novel miRNAs are given as text file.

When the relative abundances clearly indicate which the predominantly expressed miRNA is, the mature sequences are assigned names of the form, jnuhsa-233 (the predominant product) and jnuhsa-233\* (from the opposite arm of the precursor). When the data are not sufficient to determine which sequence is the predominant one, names like jnuhsa-118-5p (from the 5' arm) and jnuhsa-118-3p (from the 3' arm) have been used.

**Additional file 7 Details of the intronic novel miRNAs of the 4 samples.** List of the novel miRNAs predicted sample wise, with the information on their frequency, isomiRs, chromosomal location and prediction scores from CID-miRNA, CSHMM, miRDeep, MiPred.

**Additional file 8 Details of the intergenic novel miRNAs of the 4 samples.** List of the novel miRNAs predicted sample wise, with the information on their frequency, isomiRs, chromosomal location and prediction scores from CID-miRNA, CSHMM, miRDeep, MiPred.

**Additional file 9 A list of primers used for amplification of precursors of novel miRNAs.**

**Additional file 10 The normalized expression data along with a list of differentially regulated genes (> 1.5 fold) obtained using Ocimum microarray platform for K562 and HL60.**

## Abbreviations

miRNA: microRNA; UTR: untranslated region; RISC: RNA induced silencing complex; PTEN: phosphatase and tensin homolog; RAS: rat sarcoma viral oncogene homolog; HMGA2: high mobility group AT-hook 2; BCL2: B-cell CLL/lymphoma 2; PBMC: peripheral blood mononuclear cell; SAM: significance analysis of microarrays; FDR: false discovery rate; CML: chronic myelogenous leukemia; AML: myelogenous leukemia; ABL: c-abl oncogene 1, receptor tyrosine kinase; MEIS2: Meis homeobox 2; SMAD7: SMAD family member 7; TRIB2: tribbles homolog 2 (Drosophila); MAPK: mitogen activated protein kinase; BCL11A: B-cell CLL/lymphoma 11A (zinc finger protein); ETV4: ets variant 4; IDH1: isocitrate dehydrogenase 1 (NADP+), soluble; PPARγ: peroxisome proliferator-activated receptor gamma; EIF4A2: eukaryotic translation initiation factor 4A, isoform 2; LCK: lymphocyte-specific protein tyrosine kinase; NCOA2: nuclear receptor coactivator 2; NR4A3: nuclear receptor subfamily 4, group A, member 3; PIM1: proviral integration site 1; RNASEN: ribonuclease type III, nuclear; DGCR8: DiGeorge syndrome critical region gene 8; XPO5: exportin 5; RAN: ras-

related nuclear protein; PRMT5: protein arginine methyltransferase 5; FXR2: fragile X mental retardation, autosomal homolog 2; MOV10: Moloney leukemia virus 10; SNP: single nucleotide polymorphism

## Authors' contributions

AB and LK conceptualized the study. CV generated the computational pipeline, processed, normalized and annotated the data and did the novel miRNA prediction analysis. RK and HMA conducted the miRNA differential expression, miRNA cluster, microarray data and target prediction analyses and wet experiments. PS provided the RNA samples for sequencing. RG conducted SAM analyses. RK and AB wrote the manuscript. All the authors read and approved the final manuscript.

## Acknowledgements

The authors thank Department of Biotechnology, Government of India for their generous support and UPOE-supported HPCF for computing time. RK acknowledges support from Department of Science & Technology, GOI for a fast track grant. CV acknowledges fellowship from CSIR (CSIR SRF award no: 9/263/(0772)/9). The authors thank Dr. Narinder Sahni and Prof. Sachin Maheswari for helpful discussions. Special thanks to Chetan Aneja who helped in constructing the computational pipeline for annotating the data.

**Statement** - The authors declare no competing financial or other interest in relation to this work.

## Author Details

<sup>1</sup>School of Information Technology, Jawaharlal Nehru University, New Delhi, India, <sup>2</sup>School of Life Sciences, Jawaharlal Nehru University, New Delhi, India and <sup>3</sup>Department of Medical Oncology, Institute Rotary Cancer Hospital, All India Institute of Medical Science, New Delhi, India

Received: 8 September 2009 Accepted: 7 May 2010

Published: 7 May 2010

## References

1. Alvarez-Garcia I, Miska EA: **MicroRNA functions in animal development and human disease.** *Development* 2005, **132**:4653-4662.
2. Leung AK, Sharp PA: **microRNAs: a safeguard against turmoil?** *Cell* 2007, **130**:581-585.
3. Kulshreshtha R, Davuluri RV, Calin GA, Ivan M: **A microRNA component of the hypoxic response.** *Cell Death Differ* 2008, **15**:667-671.
4. Spizzo R, Nicoloso MS, Croce CM, Calin GA: **SnapShot: MicroRNAs in Cancer.** *Cell* 2009, **137**:586-586.e1
5. Ghildiyal M, Zamore PD: **Small silencing RNAs: an expanding universe.** *Nat Rev Genet* 2009, **10**:94-108.
6. Bartel DP: **MicroRNAs: target recognition and regulatory functions.** *Cell* 2009, **136**:215-233.
7. Kim VN, Han J, Siomi MC: **Biogenesis of small RNAs in animals.** *Nat Rev Mol Cell Biol* 2009, **10**:126-139.
8. Lee Y, Han J, Yeom KH, Jin H, Kim VN: **Drosha in primary microRNA processing.** *Cold Spring Harb Symp Quant Biol* 2006, **71**:51-57.
9. Tijsterman M, Plasterk RH: **Dicers at RISC: the mechanism of RNAi.** *Cell* 2004, **117**:1-3.
10. Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ: **miRBase: tools for microRNA genomics.** *NAR* 2008, **36**:D154-D158.
11. Rana TM: **Illuminating the silence: understanding the structure and function of small RNAs.** *Nat Rev Mol Cell Biol* 2007, **8**:23-36.
12. Hammell CM: **The microRNA-argonaute complex: a platform for mRNA modulation.** *RNA Biol* 2008, **5**:123-127.
13. Ørom UA, Nielsen FC, Lund AH: **MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation.** *Mol Cell* 2008, **30**:460-471.
14. Barbarotto E, Schmittgen TD, Calin GA: **MicroRNAs and cancer: profile, profile, profile.** *Int J Cancer* 2008, **122**:969-977.
15. Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M, Prueitt RL, Yanaihara N, Lanza G, Scarpa A, Vecchione A, Negrini M, Harris CC, Croce CM: **A microRNA expression signature of human solid tumors defines cancer gene targets.** *Proc Natl Acad Sci USA* 2006, **103**:2257-2261.
16. Deng S, Calin GA, Croce CM, Coukos G, Zhang L: **Mechanisms of microRNA deregulation in human cancer.** *Cell Cycle* 2008, **7**:2643-2646.

17. Stenvang J, Lindow M, Kauppinen S: **Targeting of microRNAs for therapeutics.** *Biochem Soc Trans* 2008, **36**:1197-1200.
18. Krichevsky AM, Gabriely G: **miR-21: a small multi-faceted RNA.** *J Cell Mol Med* 2009, **13**:39-53.
19. Meng F, Henson R, Wehbe-Janek H, Ghoshal K, Jacob ST, Patel T: **MicroRNA-21 regulates expression of the PTEN tumor suppressor gene in human hepatocellular cancer.** *Gastroenterology* 2007, **133**:647-658.
20. Takamizawa J, Konishi H, Yanagisawa K, Tomida S, Osada H, Endoh H, Harano T, Yatabe Y, Nagino M, Nimura Y, Mitsudomi T, Takahashi T: **Reduced expression of the let-7 microRNAs in human lung cancers in association with shortened postoperative survival.** *Cancer Res* 2004, **64**:3753-3756.
21. Aqeilan RI, Calin GA, Croce CM: **miR-15a and miR-16-1 in cancer: discovery, function and future perspectives.** *Cell Death Differ* 2010, **17**:215-220.
22. Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, Labourier E, Reinert KL, Brown D, Slack FJ: **RAS is regulated by the let-7 microRNA family.** *Cell* 2005, **120**:635-647.
23. Lee YS, Dutta A: **The tumor suppressor microRNA let-7 represses the HMG2 oncogene.** *Genes Dev* 2007, **21**:1025-1030.
24. Cimmino A, Calin GA, Fabbri M, Iorio MV, Ferracin M, Shimizu M, Wojcik SE, Aqeilan RI, Zupo S, Dono M, Rassenti L, Alder H, Volinia S, Liu CG, Kipps TJ, Negrini M, Croce CM: **miR-15 and miR-16 induce apoptosis by targeting BCL2.** *Proc Natl Acad Sci USA* 2005, **102**:13944-13949.
25. Yanaihara N, Caplen N, Bowman E, Seike M, Kumamoto K, Yi M, Stephens RM, Okamoto A, Yokota J, Tanaka T, Calin GA, Liu CG, Croce CM, Harris CC: **Unique microRNA molecular profiles in lung cancer diagnosis and prognosis.** *Cancer Cell* 2006, **9**:189-198.
26. Camps C, Buffa FM, Colella S, Moore J, Sotiriou C, Sheldon H, Harris AL, Gleadle JM, Ragoussis J: **hsa-miR-210 is induced by hypoxia and is an independent prognostic factor in breast cancer.** *Clin Cancer Res* 2008, **14**:1340-1348.
27. Hu X, Macdonald DM, Huettner PC, Feng Z, El Naqa IM, Schwarz JK, Mutch DG, Grigsby PW, Powell SN, Wang X: **A miR-200 microRNA cluster as prognostic marker in advanced ovarian cancer.** *Gynecol Oncol* 2009, **114**:457-464.
28. Yuan JY, Wang F, Yu J, Yang GH, Liu XL, Zhang JW: **MicroRNA-223 reversibly regulates erythroid and megakaryocytic differentiation of K562 cells.** *J Cell Mol Med* 2009, **13**:4551-4559.
29. Chivukula RR, Mendell JT: **Abate and switch: miR-145 in stem cell differentiation.** *Cell* 2009, **137**:606-608.
30. Hermeking H: **The miR-34 family in cancer and apoptosis.** *Cell Death Differ* 2010, **17**:193-199.
31. Leucht C, Stigloher C, Wizenmann A, Klafke R, Folchert A, Bally-Cuif L: **MicroRNA-9 directs late organizer activity of the midbrain-hindbrain boundary.** *Nat Neurosci* 2008, **11**:641-648.
32. Hornstein E, Mansfield JH, Yekta S, Hu JK, Harfe BD, McManus MT, Baskerville S, Bartel DP, Tabin CJ: **The microRNA miR-196 acts upstream of Hoxb8 and Shh in limb development.** *Nature* 2005, **438**:671-674.
33. Schrott GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME, Kiebler M, Greenberg ME: **A brain-specific microRNA regulates dendritic spine development.** *Nature* 2006, **439**:283-289.
34. Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, Conlon FL, Wang DZ: **The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation.** *Nat Genet* 2006, **38**:228-233.
35. Sempere LF, Freemantle S, Pitha-Rowe I, Moss E, Dmitrovsky E, Ambros V: **Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation.** *Genome Biol* 2004, **5**:R13.
36. Pfeffer S, Sewer A, Lagos-Quintana M, Sheridan R, Sander C, Grasser FA, van Dyk LF, Ho CK, Shuman S, Chien M, Russo JJ, Ju J, Randall G, Lindenbach BD, Rice CM, Simon V, Ho DD, Zavolan M, Tuschl T: **Identification of microRNAs of the herpesvirus family.** *Nat Methods* 2005, **2**:269-276.
37. Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, Nguyen JT, Barbisin M, Xu NL, Mahuvakar VR, Andersen MR, Lao KQ, Livak KJ, Guegler KJ: **Real-time quantification of microRNAs by stem-loop RT-PCR.** *Nucleic Acids Res* 2005, **33**:e179.
38. Thomson JM, Parker J, Perou CM, Hammond SM: **A custom microarray platform for analysis of microRNA gene expression.** *Nat Methods* 2004, **1**:47-53.
39. Beuvink I, Kolbf A, Budach W, Garnier A, Lange J, Natt F, Dengler U, Hall J, Filipowicz W, Weiler J: **A novel microarray approach reveals new tissue-specific signatures of known and predicted mammalian microRNAs.** *Nucleic Acids Res* 2007, **35**:e52.
40. Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Rådmark O, Kim S, Kim VN: **The nuclear RNase III Drosha initiates microRNA processing.** *Nature* 2003, **425**:415-419.
41. Lu C, Tej SS, Luo S, Haudenschild CD, Meyers BC, Green PJ: **Elucidation of the small RNA component of the transcriptome.** *Science* 2005, **309**:1567-1569.
42. Landgraf P, Rusu M, Sheridan R, Sewer A, Iovino N, Aravin A, Pfeffer S, Rice A, Kamphorst AO, Landthaler M, Lin C, Socci ND, Hermida L, Fulci V, Chiaretti S, Foà R, Schliwka J, Fuchs U, Novosel A, Müller RU, Schermer B, Bissels U, Inman J, Phan Q, Chien M, Weir DB, Choksi R, De Vita G, Frezzetti D, Trompeter HI, Hornung V, Teng G, Hartmann G, Palkovits M, Di Lauro R, Wernet P, Macino G, Rogler CE, Nagle JW, Ju J, Papavasiliou FN, Benzing T, Lichter P, Tam W, Brownstein MJ, Bosio A, Borkhardt A, Russo JJ, Sander C, Zavolan M, Tuschl T: **A mammalian microRNA expression atlas based on small RNA library sequencing.** *Cell* 2007, **129**:1401-1414.
43. Neely LA, Patel S, Garver J, Gallo M, Hackett M, McLaughlin S, Nadel M, Harris J, Gullans S, Rooke J: **A single-molecule method for the quantitation of microRNA gene expression.** *Nat Methods* 2006, **3**:41-46.
44. Moxon S, Jing R, Szittyá G, Schwach F, Rusholme Pilcher RL, Moulton V, Dalmay T: **Deep sequencing of tomato short RNAs identifies microRNAs targeting genes involved in fruit ripening.** *Genome Res* 2008, **18**:1602-1609.
45. Bar M, Wyman SK, Fritz BR, Qi J, Garg KS, Parkin RK, Kroh EM, Bendoraita A, Mitchell PS, Nelson AM, Ruzzo WL, Ware C, Radich JP, Gentleman R, Ruohola-Baker H, Tewari M: **MicroRNA discovery and profiling in human embryonic stem cells by deep sequencing of small RNA libraries.** *Stem Cells* 2008, **26**:2496-2505.
46. Glazov EA, Cottee PA, Barris WC, Moore RJ, Dalrymple BP, Tizard ML: **A microRNA catalog of the developing chicken embryo identified by a deep sequencing approach.** *Genome Res* 2008, **18**:957-964.
47. Morin RD, O'Connor MD, Griffith M, Kuchenbauer F, Delaney A, Prabhu AL, Zhao Y, McDonald H, Zeng T, Hirst M, Eaves CJ, Marra MA: **Application of massively parallel sequencing to microRNA profiling and discovery in human embryonic stem cells.** *Genome Res* 2008, **18**:610-621.
48. Friedländer MR, Chen W, Adamidi C, Maaskola J, Einspanier R, Knespel S, Rajewsky N: **Discovering microRNAs from deep sequencing data using miRDeep.** *Nat Biotechnol* 2008, **26**:407-415.
49. miRBase: the microRNA database (<http://www.mirbase.org>)
50. Liang Y, Ridzon D, Wong L, Chen C: **Characterization of microRNA expression profiles in normal human tissues.** *BMC Genomics* 2007, **8**:166.
51. Altuvia Y, Landgraf P, Lithwick G, Elefant N, Pfeffer S, Aravin A, Brownstein MJ, Tuschl T, Margalit H: **Clustering and conservation patterns of human microRNAs.** *Nucleic Acids Res* 2005, **33**:2697-2706.
52. Yu J, Wang F, Yang GH, Wang FL, Ma YN, Du ZW, Zhang JW: **Human microRNA clusters: genomic organization and expression profile in leukemia cell lines.** *Biochem Biophys Res Commun* 2006, **349**:59-68.
53. Suzuki HI, Yamagata K, Sugimoto K, Iwamoto T, Kato S, Miyazono K: **Modulation of microRNA processing by p53.** *Nature* 2009, **460**:529-533.
54. Creighton CJ, Reid JG, Gunaratne PH: **Expression profiling of microRNAs by deep sequencing.** *Brief Bioinform* 2009, **10**:490-497.
55. 't Hoen PA, Ariyurek Y, Thygesen HH, Vreugdenhil E, Vossen RH, de Menezes RX, Boer JM, van Ommen GJ, den Dunnen JT: **Deep sequencing-based expression analysis shows major advances in robustness, resolution and inter-lab portability over five microarray platforms.** *Nucleic Acids Res* 2008, **36**:e141.
56. Ro S, Park C, Young D, Sanders KM, Yan W: **Tissue-dependent paired expression of miRNAs.** *Nucleic Acids Res* 2007, **35**:5944-5953.
57. Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, Sweet-Cordero A, Ebert BL, Mak RH, Ferrando AA, Downing JR, Jacks T, Horvitz HR, Golub TR: **MicroRNA expression profiles classify human cancers.** *Nature* 2005, **435**:834-838.
58. Ramkissoon SH, Mainwaring LA, Ogasawara Y, Keyvanfar K, McCoy JP Jr, Sloan EM, Kajigaya S, Young NS: **Hematopoietic-specific microRNA expression in human cells.** *Leuk Res* 2006, **30**:643-647.
59. Agirre X, Jiménez-Velasco A, San José-Enériz E, Garate L, Bandrés E, Cordeu L, Aparicio O, Saez B, Navarro G, Vilas-Zornoza A, Pérez-Roger I, García-Foncillas J, Torres A, Heiniger A, Calasanz MJ, Fortes P, Román-Gómez J,

- Prósper F: Down-regulation of hsa-miR-10a in chronic myeloid leukemia CD34+ cells increases USF2-mediated cell growth. *Mol Cancer Res* 2008, **6**:1830-1840.
60. Venturini L, Battmer K, Castoldi M, Schultheis B, Hochhaus A, Muckenthaler MU, Ganser A, Eder M, Scherr M: Expression of the miR-17-92 polycistron in chronic myeloid leukemia (CML) CD34+ cells. *Blood* 2007, **109**:4399-4405.
61. Merkerova M, Belickova M, Bruchova H: Differential expression of microRNAs in hematopoietic cell lineages. *Eur J Haematol* 2008, **81**:304-310.
62. Genetics Home Reference [http://ghr.nlm.nih.gov/chromosome=9]
63. Hausser J, Landthaler M, Jaskiewicz L, Gaidatzis D, Zavolan M: Relative contribution of sequence and structure features to the mRNA binding of Argonaute/EIF2C-miRNA complexes and the degradation of miRNA targets. *Genome Res* 2009, **19**:2009-2020.
64. Hendrickson DG, Hogan DJ, McCullough HL, Myers JW, Herschlag D, Ferrell JE, Brown PO: Concordant regulation of translation and mRNA abundance for hundreds of targets of a human microRNA. *PLoS Biol* 2009, **7**:e1000238.
65. Xiao F, Zuo Z, Cai G, Kang S, Gao X, Li T: miRecords: an integrated resource for microRNA-target interactions. *Nucleic Acids Res* 2009:D105-110.
66. Rajewsky N: microRNA target predictions in animals. *Nat Genet* 2009, **38**(Suppl):S8-S13.
67. Fujino T, Yamazaki Y, Largaespada DA, Jenkins NA, Copeland NG, Hirokawa K, Nakamura T: Inhibition of myeloid differentiation by Hoxa9, Hoxb8, and Meis homeobox genes. *Exp Hematol* 2001, **29**:856-863.
68. Akel S, Bertolette D, Petrow-Sadowski C, Ruscetti FW: Levels of Smad7 regulate Smad and mitogen activated kinases (MAPKs) signaling and controls erythroid and megakaryocytic differentiation of erythroleukemia cells. *Platelets* 2007, **18**:566-578.
69. Stirewalt DL, Meshinchi S, Kopecky KJ, Fan W, Pogosova-Agadjanyan EL, Engel JH, Cronk MR, Dorcy KS, McQuary AR, Hockenbery D, Wood B, Heimfeld S, Radich JP: Identification of genes with abnormal expression changes in acute myeloid leukemia. *Genes Chromosomes Cancer* 2008, **47**:8-20.
70. Ruby JG, Jan CH, Bartel DP: Intronic microRNA precursors that bypass Drosha processing. *Nature* 2007, **448**:83-86.
71. Chiosea S, Jelezcova E, Chandran U, Acquaafondata M, McHale T, Sobol RW, Dhir R: Up-regulation of dicer, a component of the MicroRNA machinery, in prostate adenocarcinoma. *Am J Pathol* 2006, **169**:1812-1820.
72. Baskerville S, Bartel DP: Microarray profiling of microRNAs reveals frequent coexpression with neighboring miRNAs and host genes. *RNA* 2005, **11**:241-247.
73. Joshi T, Yan Z, Libault M, Jeong DH, Park S, Green PJ, Sherrier DJ, Farmer A, May G, Meyers BC, Xu D, Stacey G: Prediction of novel miRNAs and associated target genes in Glycine max. *BMC Bioinformatics* 2010, **11**(Suppl 1):S14.
74. Cullen BR: Transcription and processing of human microRNA precursors. *Mol Cell* 2004, **16**:861-865.
75. Johnson SM, Lin SY, Slack FJ: The time of appearance of the *C. elegans* let-7 microRNA is transcriptionally controlled utilizing a temporal regulatory element in its promoter. *Dev Biol* 2003, **259**:364-379.
76. Obernosterer G, Leuschner PJ, Alenius M, Martinez J: Post-transcriptional regulation of microRNA expression. *RNA* 2006, **12**:1161-1167.
77. Filipowicz W, Bhattacharyya SN, Sonenberg N: Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet* 2008, **9**:102-114.
78. Braun CJ, Zhang X, Savelieva I, Wolff S, Moll UM, Schepeler T, Ørntoft TF, Andersen CL, Dobbstein M: p53-Responsive microRNAs 192 and 215 are capable of inducing cell cycle arrest. *Cancer Res* 2008, **68**:10094-10104.
79. Li Z, Lu J, Sun M, Mi S, Zhang H, Luo RT, Chen P, Wang Y, Yan M, Qian Z, Neilly MB, Jin J, Zhang Y, Bohlender SK, Zhang DE, Larson RA, Le Beau MM, Thirman MJ, Golub TR, Rowley JD, Chen J: Distinct microRNA expression profiles in acute myeloid leukemia with common translocations. *Proc Natl Acad Sci USA* 2008, **105**:15535-15540.
80. Wang Y, Lee CG: MicroRNA and cancer—focus on apoptosis. *J Cell Mol Med* 2009, **13**:12-23.
81. Schoolmeesters A, Eklund T, Leake D, Vermeulen A, Smith Q, Force Aldred S, Fedorov Y: Functional profiling reveals critical role for miRNA in differentiation of human mesenchymal stem cells. *PLoS One* 2009, **4**:e5605.
82. Naguibneva I, Ameyar-Zazoua M, Poleskaya A, Ait-Si-Ali S, Groisman R, Souidi M, Cuvellier S, Harel-Bellan A: The microRNA miR-181 targets the homeobox protein Hox-A11 during mammalian myoblast differentiation. *Nat Cell Biol* 2006, **8**:278-284.
83. De Marchis ML, Ballarino M, Salvatori B, Puzzolo MC, Bozzoni I, Fatica A: A new molecular network comprising PU.1, interferon regulatory factor proteins and miR-342 stimulates ATRA-mediated granulocytic differentiation of acute promyelocytic leukemia cells. *Leukemia* 2009, **23**:856-862.
84. Lal A, Pan Y, Navarro F, Dykxhoorn DM, Moreau L, Meire E, Bentwich Z, Lieberman J, Chowdhury D: miR-24-mediated downregulation of H2AX suppresses DNA repair in terminally differentiated blood cells. *Nat Struct Mol Biol* 2009, **16**:492-498.
85. Crosby ME, Kulshreshtha R, Ivan M, Glazer PM: MicroRNA regulation of DNA repair gene expression in hypoxic stress. *Cancer Res* 2009, **69**:1221-1229.
86. Hurst DR, Edmonds MD, Scott GK, Benz CC, Vaidya KS, Welch DR: Breast cancer metastasis suppressor 1 up-regulates miR-146, which suppresses breast cancer metastasis. *Cancer Res* 2009, **69**:1279-1283.
87. Choong ML, Yang HH, McNiece I: MicroRNA expression profiling during human cord blood-derived CD34 cell erythropoiesis. *Exp Hematol* 2007, **35**:551-564.
88. Wang Q, Huang Z, Xue H, Jin C, Ju XL, Han JD, Chen YG: MicroRNA miR-24 inhibits erythropoiesis by targeting activin type I receptor ALK4. *Blood* 2008, **111**:588-595.
89. Chen CZ, Li L, Lodish HF, Bartel DP: MicroRNAs modulate hematopoietic lineage differentiation. *Science* 2004, **303**:83-86.
90. Chen CZ, Lodish HF: MicroRNAs as regulators of mammalian hematopoiesis. *Semin Immunol* 2005, **17**:155-165.
91. Wu H, Neilson JR, Kumar P, Manocha M, Shankar P, Sharp PA, Manjunath N: miRNA profiling of naive, effector and memory CD8 T cells. *PLoS One* 2007, **2**:e1020.
92. Cobb BS, Hertweck A, Smith J, O'Connor E, Graf D, Cook T, Smale ST, Sakaguchi S, Livesey FJ, Fisher AG, Merckenschlager M: A role for Dicer in immune regulation. *J Exp Med* 2006, **203**:2519-2527.
93. Gossel MJ, Hinds PW: Beyond the cell cycle: a new role for Cdk6 in differentiation. *J Cell Biochem* 2006, **97**:485-493.
94. Aghajanova L, Skottman H, Strömberg AM, Inzunza J, Lahesmaa R, Hovatta O: Expression of leukemia inhibitory factor and its receptors is increased during differentiation of human embryonic stem cells. *Fertil Steril* 2006, **86**:1193-1209.
95. Andina N, Didichenko S, Schmidt-Mende J, Dahinden CA, Simon HU: Proviral integration site for Moloney murine leukemia virus 1, but not phosphatidylinositol-3 kinase, is essential in the antiapoptotic signaling cascade initiated by IL-5 in eosinophils. *J Allergy Clin Immunol* 2009, **123**:603-611.
96. Shimizu R, Yamamoto M: Gene expression regulation and domain function of hematopoietic GATA factors. *Semin Cell Dev Biol* 2005, **16**:129-136.
97. Hu X, Li X, Valverde K, Fu X, Noguchi C, Qiu Y, Huang S: LSD1-mediated epigenetic modification is required for TAL1 function and hematopoiesis. *Proc Natl Acad Sci USA* 2009, **106**:10141-10146.
98. Bentwich I, Avniel A, Karov Y, Aharonov R, Gilad S, Barad O, Barzilai A, Einat P, Einav U, Meiri E, Sharon E, Spector Y, Bentwich Z: Identification of hundreds of conserved and nonconserved human microRNAs. *Nature Genetics* 2005, **37**:766-770.
99. Tyagi S, Vaz C, Gupta V, Bhatia R, Maheshwari S, Srinivasan A, Bhattacharya A: CID-miRNA: a web server for prediction of novel miRNA precursors in human genome. *Biochem Biophys Res Commun* 2008, **372**:831-834.
100. Agarwal S, Vaz C, Bhattacharya A, Srinivasan A: Prediction of novel precursor miRNAs using a context-sensitive hidden Markov Model (CSHMM). *BMC Bioinformatics* 2010, **11**(Suppl 1):S29.
101. Jiang P, Wu H, Wang W, Ma W, Sun X, Lu Z: MiPred: classification of real and pseudo microRNA precursors using random forest prediction model with combined features. *Nucleic Acids Res* 2007:W339-344.

doi: 10.1186/1471-2164-11-288

Cite this article as: Vaz et al.: Analysis of microRNA transcriptome by deep sequencing of small RNA libraries of peripheral blood *BMC Genomics* 2010, **11**:288