

POSTER PRESENTATION

Open Access

In vitro effect of immune regulatory cytokines on vitiligo pathogenesis

Mala Singh¹, Mohmmad Shoab Mansuri¹, Naresh C. Laddha¹, Mitesh Dwivedi¹, Yogesh S. Marfatia², Rasheedunnisa Begum^{1*}

From 2nd International Genomic Medical Conference (IGMC 2013)
Jeddah, Kingdom of Saudi Arabia. 24-27 November 2013

Background

Vitiligo is an acquired, hypomelanotic skin disorder characterized by circumscribed de-pigmented macules resulting from the loss of functional melanocytes. Various factors which may be responsible for precipitating this disorder in susceptible patients are oxidative stress, auto-immunity and neurochemicals.


Materials and methods

The skin samples were obtained with the consent of healthy individuals. Isolation of melanocytes was done according to the standard method [1] and normal human melanocytes (NHM) were grown in basal medium supplemented with growth factors. Dose dependent effect of different cytokines such as TNF α , IL6 and IL10 on NHM growth and proliferation was studied. MTT assay, RNA isolation, cDNA synthesis and relative gene expression studies were performed as described. This

study was approved by the Institutional Ethical Committee for Human Research (IECHR), The M. S. University of Baroda, Vadodara, Gujarat, India.

Results

The pro-inflammatory cytokines (TNF α and IL6) induced 37% & 20% cell death respectively in NHM, on the other hand the anti-inflammatory cytokine, IL10 did not affect the growth of NHM (Fig.1). Our earlier studies have shown high systemic mRNA and protein levels of TNF α and TNF β in Gujarat vitiligo patients compared to controls [2,3]. We have studied dose dependent effect of TNF α on NHM, and found that TNF α induced cell death in a dose dependent manner. Interestingly, higher concentrations of TNF α induced up-regulation of its receptors *TNFR1* & *TNFR2* along with significant increase in *IL6* and *ICAM1* expression (Fig. 2). IL6 was also found to increase the expression of *ICAM1*[4], which favors the attachment of



* Correspondence: rasheedunnisab@yahoo.co.in

¹Department of Biochemistry, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India
Full list of author information is available at the end of the article




Figure 2 TNF α induced expression of ICAM1, IL6, TNFR1 and TNFR2.

T cells and melanocytes and thus making the latter more susceptible for auto-immune destruction. We have also found the synergistic effect of TNF α and IL6 in inducing NHM apoptosis. In addition, TNF α and IL6 were found to aggravate their effects under oxidative stress.

Conclusions

The present study reveals that TNF α significantly induces IL6, ICAM1, TNFR1 and TNFR2 expression. In addition IL6 also induces ICAM1 expression [4]. ICAM1 enhances T-cell and melanocyte attachment, thus augmenting melanocyte destruction by immune system. Under oxidative stress, which mimics the microenvironment of vitiligo, TNFA is found to enhance apoptosis of melanocytes which would result in de-pigmentation of the skin. Thus, our *in vitro* studies further strengthen the scientific evidences linking oxidative stress and immune system to vitiligo pathogenesis giving credence to a convergent terminal pathway of oxidative stress-autoimmunity mediated melanocyte loss.

Authors' details

¹Department of Biochemistry, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India. ²Department of Skin and V.D., Faculty of Medicine, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India.

Published: 2 April 2014

References

1. Czajkowski R, Placek W, Drewa T, Kowalyszyn B, Sir J, Weiss W: Autologous cultured melanocytes in vitiligo treatment. *Dermatol Surg* 2007, **33**:1027-1036.
2. Laddha NC, Dwivedi M, Begum R: Increased Tumor Necrosis Factor (TNF)- α and its promoter polymorphisms correlate with disease progression and higher susceptibility towards vitiligo. *PLoS ONE* 2012, **7**:e52298.
3. Laddha NC, Dwivedi M, Gani AR, Mansuri MS, Begum R: Tumor Necrosis Factor B (TNFB) genetic variants and its increased expression are associated with vitiligo susceptibility. *PLoS ONE* 2013, **8**:e81736.
4. Wung BS, Ni CW, Wang DL: ICAM-1 induction by TNF alpha and IL-6 is mediated by distinct pathways via Rac in endothelial cells. *J Biomed Sci* 2005, **12**:91-101.

doi:10.1186/1471-2164-15-S2-P39

Cite this article as: Singh et al.: *In vitro* effect of immune regulatory cytokines on vitiligo pathogenesis. *BMC Genomics* 2014 15(Suppl 2):P39.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 BioMed Central