

RESEARCH ARTICLE

Open Access



# Genomic analyses of aminoacyl tRNA synthetases from human-infecting helminths

Preeti Goel<sup>1,2</sup>, Suhel Parvez<sup>2</sup> and Amit Sharma<sup>1\*</sup>

## Abstract

**Background:** Helminth infections affect ~ 60% of the human population that lives in tropical and subtropical regions worldwide. These infections result in diseases like schistosomiasis, lymphatic filariasis, river blindness and echinococcosis. Here we provide a comprehensive computational analysis of the aminoacyl tRNA synthetase (aaRS) enzyme family from 27 human-infecting helminths. Our analyses support the idea that several helminth aaRSs can be targeted for drug repurposing or for development of new drugs. For experimental validation, we focused on Onchocerciasis (also known as “river blindness”), a filarial vector-borne disease that is prevalent in Africa and Latin America. We show that halofuginone (HF) can act as a potent inhibitor of *Onchocerca volvulus* prolyl tRNA synthetase (OvPRS).

**Results:** The conserved enzyme family of aaRSs has been validated as druggable targets in numerous eukaryotic parasites. We thus embarked on assessing aaRSs from the genomes of 27 helminths that cause infections in humans. In order to delineate the distribution of aaRSs per genome we utilized Hidden Markov Models of aaRS catalytic domains to identify all orthologues. We note that *Fasciola hepatica* genome encodes the highest number of aaRS-like proteins (69) whereas *Taenia asiatica* has the lowest count (32). The number of genes for any particular aaRS-like protein varies from 1 to 8 in these 27 studied helminths. Sequence alignments of helminth-encoded lysyl, prolyl, leucyl and threonyl tRNA synthetases suggest that various known aaRS inhibitors like Cladosporin, Halofuginone, Benzoborale and Borrelidin may be of utility against helminths. The recombinantly expressed *Onchocerca volvulus* PRS was used as proof of concept for targeting aaRS with drug-like molecules like HF.

**Conclusions:** Systematic analysis of unique subdomains within helminth aaRSs reveals the presence of a number of non-canonical domains like PAC3, Utp-14, Pex2\_Pex12 fused to catalytic domains in the predicted helminth aaRSs. We have established a platform for biochemical validation of a large number of helminth aaRSs that can be targeted using available inhibitors to jump-start drug repurposing against human helminths.

**Keywords:** Aminoacyl tRNA synthetases, Cladosporin, Halofuginone, Helminth, Inhibitors

## Background

Helminths are common infectious agents of humans in developing countries and cause significant mortality and morbidity [1]. Helminths are comprised of nematodes (roundworms- intestinal and filarial worms) as well as of platyhelminths (flatworms- flukes and schistosomes) [1]. Prevalence of soil transmitted helminth infections remains high in Sub-Saharan Africa, Latin America and

Asia [2]. Helminth infections also contribute to physical disabilities, anemia, malnourishment and result in decreased productivity in the workplace [1, 3]. Some common diseases caused by helminths are river blindness, echinococcosis, dracunculiasis, taeniasis, schistosomiasis and lymphatic filariasis. Schistosomiasis alone infects ~ 230 million people in the developing world [4]. Both Hookworm and Schistosomiasis threaten pregnant women as these infections increase the risk of child and maternal morbidity [5]. Among people living in impoverished areas of developing countries, Onchocerciasis is prevalent and causes skin disease and visual impairment,

\* Correspondence: amitsharma68@gmail.com

<sup>1</sup>Structural Parasitology Group, International Centre for Genetic Engineering and Biotechnology (ICGEB), New Delhi 110067, India

Full list of author information is available at the end of the article



while limb and genital deformities are the outcome of lymphatic filariasis [1]. Taeniasis caused by *Taenia solium* is known to cause blindness, headache, convulsions and epileptic seizures that can be fatal [6]. The threadworm *Strongyloides stercoralis* is found in digestive tracts where it can damage the duodenum and produce abdominal pain, epigastric tenderness and diarrhea [7]. Thus, helminth infections are a serious cause of poor health in many countries.

Despite the success of MDA (Mass Drug Administration) programmes, helminth diseases have not been eliminated in endemic areas and limitations have been observed in the effectiveness of treatments [1, 8, 9]. A list of known anthelmintics with their mechanism of action is provided in Table 1. There is concern of drug resistance to prime drugs including Benzimidazoles and Ivermectin [10–12]. Ivermectin is the drug of choice in helminth infections but is contraindicated in high-intensity *Loa loa* infection due to potential fatal side-effects [13, 14]. For soil-transmitted helminth infections, the most widely used drugs Albendazole and Mebendazole have shortcomings in their efficacy profiles, especially against infections by hookworm and *T. trichiura* [15, 16]. Hence, there is a need for the discovery of new druggable targets and inhibitor scaffolds in the context of human helminth infections.

Protein synthesis machinery is an important focus of drug targeting against many human pathogens. The aaRSs constitute a family of enzymes that are responsible for transfer of amino acid to their cognate tRNA molecules, a step integral

to synthesis of proteins. The aaRSs are an important component of protein translational machinery in all cells, and have been validated as druggable targets. Few examples of aaRSs inhibitors include Cladosporin (CLD), Halofuginone (HF), Borrelidin and Benzoborale AN2690 (5-fluoro-1, 3-dihydro-1-hydroxy-2, 1-benzoxaborole) [17]. CLD competitively inhibits malaria parasite lysyl tRNA synthetase (KRS). It mimics ribose and adenine moieties of ATP as it has a THP ring (2, 6-disubstituted tetrahydropyran) fused to an iso-coumarin moiety [18]. HF has a quinazolinone ring fused to hydroxypiperidine ring and these respectively mimic the tRNA adenine and L-proline. HF is a potential inhibitor of prolyl tRNA synthetase of *Plasmodium falciparum* [19]. Borrelidin is known to inhibit threonyl tRNA synthetase (TRS). Borrelidin is an 18-membered macrolide-polyketide which simultaneously binds to four distinct pockets of ATP, amino acid, tRNA binding site and a fourth accessory site [20]. Benzoborale AN2690 (5-fluoro-1, 3-dihydro-1-hydroxy-2, 1-benzoxaborole) is known to block leucyl tRNA synthetase by covalently trapping tRNA in the editing site of the enzyme [21]. Apart from their translational functions, aaRSs are associated with various non-canonical functions such as role in angiogenesis, apoptosis, inflammation and immune responses [22]. Paralogs of aaRSs exist that have lost their aminoacylation functions and have adapted other roles [22, 23]. Hence our computational work lays the foundation for experimental dissection of helminth aaRSs and for understanding roles of aaRSs in parasite biology.

**Table 1** Major anthelmintic drugs and their mechanisms of action

Anthelmintic drugs	Effective against	Mechanism of action
Ivermectin	Filariasis, Onchocerciasis	Targets GABA receptors of parasite, chloride ion influx enhances hyperpolarisation and causes paralysis of worm
Benzimidazoles- Albendazole Mebendazole Thiabendazole	Threadworm, Hookworm, Roundworms	Binds to beta-tubulin and prevents its polymerisation
Praiquantel	Tapeworm, Schistosomiasis	Increase Ca <sup>2+</sup> permeability, causing muscle paralysis
Piperazine	Threadworm, Roundworm	Weak GABA mimetic and cause flaccid
Niclosamide	Tapeworm	Inhibits ATP generation by preventing phosphorylation of ADP in mitochondria
Diethylcarbamazine	Filariasis,Loiasis, Onchocerciasis	Interferes with cyclooxygenase and lipooxygenase pathway reducing eicosanoid production
Pyrantel pamoate Levamisole Morantel	Most Nematode infections	Nicotinic receptor agonist, elicit spastic muscle paralysis
Paraherquamide	Nematodes	It is a competitive inhibitor of the body wall nACh receptor, causes flaccid paralysis

**Results**

**Phylogenetic analyses of human-infecting helminth**


Humans are parasitized by two major groups of parasitic worms - the Nematoda (roundworms) and Platyhelminthes (flatworms). Platyhelminthes are part of the superphylum Lophotrochozoa whereas Nematoda is part of the superphylum Ecdysozoa - these two phyla are not related zoologically. Trematoda and Cestoda are likely monophyletic and have different pre-turbellarian ancestors. Trematoda (endoparasitic flukes) and Cestoda (endoparasitic tapeworms) have evolved in different directions each having unique morphology and life style [24, 25]. We constructed phylogenetic tree of the 27 human-infecting helminths based on their whole proteomes (Fig. 1). The tree displays groups of helminths as platyhelminths and nemathelminths.

The nematodes (round worms) are major intestinal worms and platyhelminths include cestodes (tapeworms) and trematodes (blood, tissue and organ flukes).

Phylogenetic analyses reveal close evolutionary linkage between various human-infecting helminths. However,

helminths belonging to the same phylogenetic group (trematodes, cestodes and nematodes) do not cause similar human infections. Therefore above three groups are not divided based on types of infection caused or prevalence in any particular region.

To study the evolutionary aspects of aminoacyl tRNA synthetases present in these 27 helminths, as an example, we studied the evolutionary linkages of the two PRSs in *Onchocerca volvulus* of its total of 39 aaRSs. All *Onchocerca volvulus* aaRSs were evaluated for the presence of signal peptides and predictions suggest that 9 out of the 39 are likely to be mitochondrial. We surmised that of the two PRSs in *Onchocerca volvulus* one is likely cytoplasmic. We subjected both PRSs to phylogenetic analyses in order to determine their evolutionary relatedness to bacterial and eukaryotic aaRSs [26]. Interestingly, one OvPRS gene (OVOC 1179) clustered with bacterial PRSs and with mitochondrial PRSs in eukaryotes whereas the other OvPRS (OVOC 11842) clustered with cytoplasmic eukaryotic PRSs indicating different evolutionary origins for the two *Onchocerca volvulus*



**Fig. 1** Phylogenetic tree of human infecting helminths based on their proteomes. Pink circle shows Trematodes, green shows Cestodes and the blue shows Nematodes

PRs (Fig. 7). This phylogenetic analysis thus can be extended to annotate the evolutionary relatedness of all helminth aaRSs with each other. It is thus likely that the cytoplasmic and mitochondrial versions of aaRSs in these helminths have eukaryotic-like and prokaryotic-like lineages.

#### Aminoacyl tRNA synthetase-like proteins in helminths

Aminoacyl tRNA synthetases are important enzymes of protein translation machinery and are also involved in various non canonical functions. It has been fruitful to compile genome-wide data on the distribution of aaRSs in various pathogens [27–29]. Comparative analyses of aaRSs genes has revealed that numerous genomes contain open reading frames that encode synthetase-like proteins that have a catalytic domain which is specific for a given tRNA synthetase [27]. Amongst these aaRS-like proteins, many are smaller than the corresponding functional tRNA synthetases [17]. It seems that evolution has recruited this family of enzymes into a range of diverse non-canonical, non aminoacylation functions [17]. Using similar ideas, we catalogued all aaRS-like proteins in the 27 human infecting helminths. The proteins with tRNA synthetase catalytic domains were identified based on HMM-profile searches, and then categorized into specific aaRS-like proteins based on blast searches against a database of well-annotated aaRSs (AARS database). Predominantly, an aaRS consists of a catalytic domain (which is involved in aminoacylation) and an anticodon binding domain (which binds to the anticodon region of the tRNA and ensures binding of tRNA to the correct amino acid). Some aaRSs have editing domains that cleave incorrectly paired amino acyl-tRNA molecules and few additional non canonical domains are also present in some tRNA synthetases [29, 30].

Interestingly different numbers of aaRS-like proteins are predicted in the 27 helminth genomes (Fig. 2 and Additional file 1: Table S1). For example, there are 69 aaRS-like proteins in *Fasciola hepatica* and this may be related to its large genome size. For regular protein synthesis, the expectation would be of 20 aaRSs each for cytoplasmic and mitochondrial translation, however, this does not hold true for many of the helminths (Fig. 2). We also observed multiple aaRS-like proteins for any particular amino acid in several helminth genomes - for example 8 NRSs in *Brugia malayi* and 7 DRSs in *Schistosoma mansoni* (Fig. 3). In *Brugia malayi*, a 5 gene cluster of NRS (IDs: Bm17689, Bm9833, Bm11983, Bm3517 and Bm749) is present on chromosome 3 (Fig. 3). These 5 genes encode for cytoplasmic NRS in this organism and this protein also serves as an antigenic determinant in *Brugia malayi* [31]. The cytoplasmic NRSs in *Brugia*


*malayi* are secreted from the helminth and these NRS genes encode an immunodominant antigen which is responsible for strong antibody response in humans with lymphatic filariasis [31]. The atomic structure of *B. malayi* NRS suggests two structured domains: (1) an amino-terminal domain in which 80 amino acids fold as for interleukin-8 (IL-8) and interacts with the extracellular loops of G-protein coupled IL-8 receptors and (2) a 438 amino acid catalytic domain [32]. It has been shown that this *BmNRS* promotes pro-inflammatory activities such as chemotaxis of cells that express IL-8 receptors [33]. The *Brugia malayi* NRS is an example of aaRSs that are present in high copy number and that likely perform additional non-canonical functions.

In *Schistosoma mansoni*, 7 DRS-like proteins were found and each of the respective genes vary in sequence and chromosomal location (Fig. 3). Transcriptome data indicates expression of 4 of 7 DRS-like genes in *Schistosoma mansoni* [34]. However, it remains unclear if all of these genes encode functional aaRSs although they encode catalytic domains of different molecular sizes (Fig. 3). In general, we found that several aaRS-like proteins such as alanyl, aspartyl, asparaginyl and isoleucyl aaRSs are present in multiple copies within the helminth genomes whereas threonyl, cysteinyl, glycyl and histidyl aaRSs are present as a single copy (Fig. 2). Apart from the presence of an unusually large number of aaRS-like proteins (total of 38) in *Schistosoma mansoni*, we were intrigued to note that no discernible TRS was identified in our searches. However, in literature two putative threonyl tRNA synthetases have been reported in *Schistosoma mansoni* [35], but we note that neither of these have the canonical catalytic domain of a tRNA synthetase. Additionally, our searches failed to find TRS and GRS in *Schistosoma japonicum*.

We also found numerous non-canonical domains fused with the catalytic domains in helminth aaRS-like proteins (Additional file 2: Table S2). Two interesting examples are: (1) presence of PAC3 (proteasome assembly chaperone 3 proteins) domain in IRSs from *Brugia malayi* and *Necator americanus*, (2) presence of Utp-14 (part of a large ribonucleoprotein protein complex containing the U3 snoRNA) in PRs from *Ascaris lumbricoides*, *Loa loa* and *Dracunculus medinensis* (Fig. 4). A complete list of such unusual domain fusions with helminth aaRSs is provided in Additional file 2: Table S2.

#### Sequence and structural similarities in aaRS-like proteins

We next decided to focus on specific tRNA synthetases in an effort to study their druggable properties. Analysis of lysyl tRNA synthetase from *Onchocerca volvulus* suggests the presence of only one copy of lysyl tRNA synthetase (KRS, gene ID- Ov2569) in it. We assessed via




sequence alignments the possible structural similarities between *Onchocerca volvulus* KRS and *Plasmodium falciparum* KRS where the drug Cladosporin binds (Fig. 5b) [18, 35, 36]. We noted considerable sequence identity between *Ov/Pf* KRS cladosporin interacting residues (PDB ID: 4PG3). Similar analysis was done for *Brugia malayi* TRS (gene ID - Bm13920) using the complex *E. coli* TRS with the drug Borrelidin (PDB ID 4P3O). There are two LRSs predicted in *Brugia malayi* (Bm5408 and Bm7489). Sequence alignments with LRS of *Cryptosporidium muris* (PDB ID 5FOM) show conservation of benzoborale binding residues in one of the predicted LRS of *Brugia malayi* (Bm5408). There are three predicted PRSs (Gene IDs Bm11475, Bm6054, Bm5875) in *Brugia malayi* and in other helminths like *Onchocerca volvulus*. Sequence alignments of these with *Pf*PRS show conservation in HF interacting residues for many helminths. These sequence and structural comparisons suggest that known inhibitors of tRNA synthetases may be of utility as

starting points of drug discovery against helminths. Assessment of sequence conservation within inhibitor-binding residues in enzyme-inhibitor complexes can be used to identify potential lead compounds and their molecular targets in multiple helminths (Fig. 5). We then decided to validate these observations by recombinantly producing the *Ov*PRS.

**Activity and inhibition by halofuginone**

Halofuginone is a known inhibitor of *Plasmodium falciparum* PRS [19]. HF was tested against *Onchocerca volvulus* PRS based on the binding site sequence similarities between *Pf*PRS and *Ov*PRS. Assay conditions were optimized to determine the activity of *Ov*PRS in absence and presence of HF (Fig.6). This involved determining the optimal enzyme concentration, the linearity of assay with respect to time, and the Michaelis Menten constants for proline and ATP (Fig. 6 a-e). The apparent  $K_m$  for proline and ATP were determined to be  $24.17 \pm 11 \mu M$  and  $18.9 \pm 2.2 \mu M$ .



Interestingly, HF showed an IC<sub>50</sub> value of ~ 30 nM against OvPRS (Fig. 6f). These results thus highlight the utility of HF against OvPRS as also potentially against other helminth PRSs.


## Discussion

Treatment and prevention of infections caused by helminths presents an emerging challenge. There is an imminent need to strengthen efforts to identify new anti-parasitic agents with alternative modes of action against helminth diseases. Dissecting and understanding the critical components of helminth protein production machinery will help establish a platform for targeting specific helminth aaRSs. Data from several laboratories over the past decade has shown that the aaRS family of enzymes from eukaryotic parasites are druggable [18, 19, 37–42]. The human/pathogen aaRSs can at times show minor/major differences in their active site residues or additional residues thereby conferring selectivity [36]. The idea of STOPP (Structure-based targeting of orthologous pathogen proteins) has been implemented for pathogen aaRSs [41]. Two examples of STOPP are (1) halofuginone in context of PRSs from *Plasmodium falciparum*, *Toxoplasma gondii*, *Cryptosporidium*

*parvum*, *Leishmania major* and *Eimeria tenella* [43], and (2) cladospirin in context of KRSs of *Plasmodium falciparum*, *Loa loa* and *Schistosoma mansoni* [35].

In this work, we have studied the distribution of aaRS-like proteins in 27 human infecting helminths. The number of any particular aminoacyl tRNA synthetase varies from 1 to 8 in the studied group of helminths. Ideally there should be only two copies (one cytoplasmic and one mitochondrial) for a particular aaRS in each helminth but this does not hold true for most of the 27 studied organisms as shown in Fig. 2.

We have modelled four aminoacyl tRNA synthetases from *Brugia malayi* and *Onchocerca volvulus* in order to investigate whether these helminth enzymes may be druggable just as their homologs are [19, 37–39]. The currently known anti-aaRSs compounds can be utilised to target these four helminth aaRSs and to jumpstart drug repurposing. We analysed multiple sequence alignments to assess the binding sites within aaRSs (PRS, KRS, LRS and TRS) for the following inhibitors: halofuginone, cladospirin, benzoborole and borellidin in the 27 studied human-infecting helminths (Additional file 3: Figure S1). Several helminth aaRSs display high conservation in the above binding residues and indeed



modelling presented here shows the feasibility of drug targeting.


## Conclusions

There is an immediate need to find new targets and drugs for helminth infections due to the potential of evolving drug resistance. The novelty of this study lies in computational analysis that combines Hidden Markov Models (HMM) and similarity searches against tRNA synthetase catalytic domains to identify all aaRSs in 27 human-infecting helminths. This is the first report of an extensive analysis of helminth aaRSs, and sets up avenues for experimental validation of our observations. If pursued, helminth aaRSs may prove to be of utility as novel targets for anti-helminth drug development.

## Methods

### Data collection and computational analysis


Whole proteome FASTA files for human-infecting helminths were downloaded from WormBase parasite database [44]. HMMER software package and HMMs for catalytic domains of tRNA synthetases were downloaded from pfam database [45]. To identify potential aaRSs, hmm searches were carried out using tRNA synthetases catalytic domains (tRNA-synt\_1b, tRNA-synt\_1f, tRNA-synt\_2, tRNA-synt\_2b, tRNA-synt\_2c, tRNA-synt\_1e, tRNA-synt\_1g, tRNA-synt\_2d, tRNA-synt\_1d, tRNA-synt\_1, tRNA-synt\_2e, tRNA-synt\_1c) with gathering threshold. This threshold is the default setting for hmmscan which instructs HMMER to use the sequence and hit thresholds defined in the searched HMM files. These values are set conservatively to ensure



that there are no known false positives. These searches identified 1071 hits in 27 helminth genomes. The identified hits were annotated to specific tRNA synthetase by using BLAST searches against annotated tRNA synthetases downloaded from AARSs database using e-value of  $1e^{-6}$  [46]. Total of 1070 (Additional file 1: Table S1) of 1071 hits identified from hmm searches also gave at least 10 significant hits in the BLAST searches against annotated tRNA synthetase from AARS DataBase. Any given tRNA synthetase was annotated to specific amino acid if majority of the top 10 significant BLAST hits were for particular amino acid. Non-canonical domains in aaRSs were identified by analyzing hmm searches of all domains in Pfam-A database using gathering threshold. Orthologs of human infecting helminths were identified using the best bidirectional hit algorithm using BLASTP with  $1e^{-6}$  as the e-value cutoff [47]. There were 687 core genes in these 27 genomes with orthologs which were then aligned using clustalO [48]. The alignments were cleaned using

GBlocks with default parameters to remove non-informative and gapped sites [49]. A maximum likelihood phylogenetic tree was constructed using RAxML with 100 bootstraps [50, 51]. The substitution model was automatically determined by RAxML. The tree was visualized using dendroscope [52]. Sequences of the two OvPRSs (OVOC1179 AND OVOC11842) were retrieved from WormBase parasite database and a BLAST search was carried against them. The resulting sequences from different organisms were then aligned using Clustalw and a phylogenetic tree was constructed in MEGA7 [53]. The evolutionary history was inferred by using Maximum Likelihood method based on the JTT matrix-based model. The tree with highest log likelihood (-6518.26) is shown in (Fig. 7) Initial tree (s) for the heuristic searches were obtained automatically by applying Neighbour-Join and BioNJ algorithms to a matrix of pairwise distances estimated using a JTT model, we then selected the topology with superior log likelihood value. The tree is





**Fig. 6** Enzymatic assays of *Onchocerca volvulus* PRS. **a)** Purified OvPRS on SDS-PAGE **b)** Aminoacylation assay using malachite green dye (53). Linearity of the assay with respect to enzyme concentration. **c)** Assay linearity with respect to time, red shows control without enzyme and the blue is reaction in presence of enzyme. **d, e)** Proline Km determination in the presence of a saturating concentration of ATP (200  $\mu$ M), ATP Km determination in the presence of saturating concentration of proline (1 mM). **f)** Enzyme inhibition with halofuginone using 50  $\mu$ M proline, 150  $\mu$ M ATP and 150 nM OvPRS enzyme. The IC<sub>50</sub> of halofuginone for inhibition of OvPRS is estimated to be  $\sim$  30 nM


drawn to scale, with branch lengths measured in the number of substitutions per site. All positions containing gaps and missing data were eliminated. The analysis involved 22 amino acid sequences. Evolutionary analyses were conducted in MEGA7.

Transcriptome data for *Schistosoma mansoni* were obtained from Lu Z et al. [34] Protein sequences of aaRSs (lysyl, leucyl, prolyl, and threonyl) and their structural information were downloaded from databases. A multiple alignment study was performed on lysyl, leucyl, prolyl and threonyl tRNA synthetases of human helminths using clustal Omega. Drug binding residues for each tRNA synthetase were identified from structure and literature. The structures of leucyl and threonyl tRNA synthetases of *Brugia malayi* were modelled and compared with the structure of already known leucyl

tRNA synthetase of *Cryptosporidium muris* (PDB ID 5FOM) and threonyl tRNA synthetase of *E.coli* respectively (PDB ID4P3O). Lysyl and prolyl tRNA synthetase of *Onchocerca volvulus* were modelled and compared to lysyl and prolyl tRNA synthetases from *Plasmodium falciparum* (PDB IDs: 4YCV, 4YDQ).

#### Protein expression and purification of OvPRS

The OvPRS sequence was retrieved from the WormBase database. We made a smaller construct of full-length OvPRS on the basis of structural modelling. The construct was designed for optimal expression in *E. coli*. The OvPRS construct consists of 497 amino acids (279–775; 57 kDa). This construct was sub-cloned in pETM-41 (maltose binding protein with histidine tag). Recombinant OvPRS was produced in *E. coli* BL21



codon+ cells. Protein expression was induced by adding 0.5 mM isopropyl b-D-thiogalactopyranoside (IPTG) to cells grown at 37 °C for 4 h, and for 20 h post-induction at 18 °C. Cells were obtained by centrifugation at 5000 g for 15 min. Bacterial cell pellet was resuspended in a buffer containing 50 mM Tris–HCl pH 8.0, 500 mM NaCl, 10 mM bME, 15% v/v glycerol, 0.1 mg ml<sup>-1</sup> lysozyme and EDTA free protease inhibitor cocktail (Roche). Cells were lysed by sonication and cleared by centrifugation at 13,000 g for 45 min. The supernatant was applied to amylose beads (NEB) and protein was eluted using the buffer (50 mM Tris–HCl pH 8.0, 250 mM NaCl, 10 mM bME, 10 mM maltose). TEV protease was added and kept for 24 h at 20 °C to remove the tag. Cleaved OvPRs protein was purified by gel filtration chromatography on Superdex 200 column 16/60 GL (GE Healthcare) which was equilibrated with 20 mM Tris-HCl pH 8, 250 mM NaCl and 2 mM DTT. Eluted OvPRs fractions were observed on SDS-PAGE and the pure ones were pooled, concentrated to 15 mg/ml and stored at – 80 °C.

#### Aminoacylation assays

The aminoacylation reactions (100 µl total volume each) were performed in clear, flat-bottomed, 96-well plates (Costar 96-well standard microplates), and the reaction mixtures were incubated for 1 h at 37 °C. In the first step of amino acylation reaction pyrophosphate is released which is converted to inorganic phosphate by PPIase. This liberated inorganic phosphate was detected by addition of 25 µl of malachite

green solution and incubated for 10 min at room temperature. Malachite green is a dye that detects the PPI formed in the reaction which correlates with enzyme activity [54]. This method is based on the formation of a complex between malachite green molybdate and free inorganic phosphates that absorbs at 620–640 nm. Absorbance was measured at 620 nm using a spectrophotometer. Reactions without enzyme or without L-Pro were performed as background controls, and data from reactions without enzyme were subtracted from the measurements. 10 mM EDTA was added to the reaction mixture at different time points (5, 10, 20, 30, 40–180 min) and mixed on ice to stop the reaction in order to observe the dependence of enzyme activity on time. For OvPRs inhibition, a reaction solution containing PRs (0.15 µM) was mixed with HF at 100 µM–0.1 nM (10 fold dilutions).

#### Additional files

**Additional file 1: Table S1.** A list of all aaRSs like proteins from 27 studied human infecting helminths along with their gene IDs. (PDF 97 kb)

**Additional file 2: Table S2.** Domain description of all the aaRSs like proteins showing the presence of unusual domains. Unusual domains are mentioned in red. (PDF 99 kb)

**Additional file 3: Figure S1.** Multiple sequence alignment (MSA) of aaRSs 1) PRs of *Plasmodium falciparum* with PRs of all studied human-infecting helminths showing conservation in HF binding residues, 2) CLD binding residues in *Plasmodium falciparum* aligned with KRSs of helminths, 3) binding residues in TRS of *Cryptosporidium muris* aligned with

TRSs of helminths, 4) Benzoborale binding residues in LRS of *E.coli* aligned with LRSs of helminths. (DOCX 1888 kb)

### Abbreviations

aaRSs: Aminoacyl tRNA synthetases; AMP: Adenosine mono phosphate; ATP: Adenosine tri phosphate; CLD: Cladosporin; DRS: Aspartyl tRNA synthetase; HF: Halofuginone; HMM: Hidden markov model; IC50: Inhibitory concentration 50; IL-8: Interleukin-8; Km: Michaelis Menten constant; KRS: Lysyl tRNA synthetase; LRS: Leucyl tRNA synthetase; MDA: Mass Drug Administration; NRS: Asparaginyl tRNA synthetase; OvPRS: *Onchocerca volvulus* prolyl tRNA synthetase; PAC3: Proteasome assembly chaperone 3; THP: 2,6-disubstituted tetrahydropyran; TRS: Threonyl tRNA synthetase

### Acknowledgements

We thank Dr. Rai for constant encouragement and Dr. GD Rekoj for insights.

### Funding

AS laboratory is supported by MMV, GHIT, DST, DBT and COE grants. PG is supported by UGC fellowship. The funding in our laboratory did not influence the study design or its analysis.

### Availability of data and materials

The data utilised are as follows  
The genomes of 27 studied helminths are taken from WormBase parasite database (<https://parasite.wormbase.org/index.html>). For blast studies AARS database is utilised which is a compilation of amino acid sequences of all known AARSs, which are available at <http://biobases.ibch.poznan.pl/aars/>. For HMM model and domain analyses pfam database is used (<https://pfam.xfam.org/>). All other data generated is provided in the Additional files.

### Authors' contributions

PG, SP and AS designed the study and wrote the manuscript. All authors have read and approved the manuscript.

### Ethics approval and consent to participate

Not applicable for this study.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

### Author details

<sup>1</sup>Structural Parasitology Group, International Centre for Genetic Engineering and Biotechnology (ICGEB), New Delhi 110067, India. <sup>2</sup>Department of Toxicology, School of Chemical and Life Sciences, Jamia Hamdard, New Delhi 110063, India.

Received: 21 December 2018 Accepted: 9 April 2019

Published online: 02 May 2019

### References

- Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth infections: the great neglected tropical diseases. *J Clin Invest*. 2008;118:1311–21.
- Pullan RL, Smith JL, Jasrasaria R, Brooker SJ. Global numbers of infection and disease burden of soil-transmitted helminth infections in 2010. *Parasit Vectors*. 2014;7:1–19.
- Hotez PJ, Ferris MT. The antipoverty vaccines. *Vaccine*. 2006;24:5787–99.
- Colley DG, Bustinduy AL, Secor WE, King CH. Human schistosomiasis. *Lancet*. 2014;383:2253–64 Elsevier Ltd.
- Christian P, Khatry SK, West PKP. Antenatal anthelmintic treatment, birthweight, and infant survival in rural Nepal. *Lancet*. 2004;364:981–3.
- García HH, Gonzalez AE, Evans CA, Gilman RH. Cysticercosis working Group in Peru Taenia solium cysticercosis. *Lancet*. 2003;362(9383):547–56.
- Mayayo E, Gomez-Aracil V, Azua-Blanco J, Azua-Romeo J, Capilla J, Mayayo R. Strongyloides stercoralis infection mimicking a malignant tumour in a non-immunocompromised patient. Diagnosis by bronchoalveolar cytology. *J Clin Pathol*. 2005;58:420–2.
- Flohr C, Tuyen LN, Lewis S, Minh TT, Campbell J, Britton J, et al. Low efficacy of mebendazole against hookworm in Vietnam: two randomized controlled trials. *Am J Trop Med Hyg*. 2007;76:732–6.
- Geerts S, Gryseels B. Drug resistance in human helminths: current situation and lessons from livestock. *Clin Microbiol Rev*. 2000;13:207–22.
- Albonico M, Engels D, Savioli L. Monitoring drug efficacy and early detection of drug resistance in human soil-transmitted nematodes: a pressing public health agenda for helminth control. *Int J Parasitol*. 2004;34:1205–10.
- Osei-Atweneboana MY, Eng JK, Boakye DA, Gyapong JO, Prichard RK. Prevalence and intensity of *Onchocerca volvulus* infection and efficacy of ivermectin in endemic communities in Ghana: a two-phase epidemiological study. *Lancet*. 2007;369:2021–9.
- Bourguinat C, Pion SDS, Kamgno J, Gardon J, Duke BOL, Boussinesq M, et al. Genetic selection of low fertile *Onchocerca volvulus* by ivermectin treatment. *PLoS Negl Trop Dis*. 2007;1(1):e72.
- Chippaux JP, Boussinesq M, Gardon J, Gardon-Wendel N, Ernould JC. Severe adverse reaction risks during mass treatment with ivermectin in loiasis-endemic areas. *Parasitol Today*. 1996;12:448–50.
- Gardon J, Gardon-Wendel N, Demanga-Ngangue KJ, Chippaux JP, Boussinesq M. Serious reactions after mass treatment of onchocerciasis with ivermectin in an area endemic for Loa loa infection. *Lancet*. 1997;350:18–22.
- Moser W, Schindler C, Keiser J. Efficacy of recommended drugs against soil-transmitted helminths: systematic review and network meta-analysis. *BMJ*. 2017;358:1–10.
- Veracruzse J, Albonico M, Behnke JM, Kotze AC, Prichard RK, McCarthy JS, et al. Is anthelmintic resistance a concern for the control of human soil-transmitted helminths? *Int J Parasitol Drugs Drug Resist*. 2011;1:14–27 Australian Society for Parasitology.
- Manickam Y, Chaturvedi R, Babbar P, Malhotra N, Jain V, Sharma A. Drug targeting of one or more aminoacyl-tRNA synthetase in the malaria parasite *Plasmodium falciparum*. *Drug Discov Today*. 2018;23(6):1233–40.
- Sharma A, Khan S, Sharma A, Belrhali H, Yogavel M. Structural basis of malaria parasite lysyl-tRNA synthetase inhibition by cladosporin. *J Struct Funct Genom*. 2014;15:63–71.
- Jain V, Yogavel M, Oshima Y, Kikuchi H, Touquet B, Hakimi MA, et al. Structure of prolyl-tRNA synthetase-halofuginone complex provides basis for development of drugs against malaria and toxoplasmosis. *Structure*. 2015;23(5):819–29.
- Fang P, Yu X, Jeong SJ, Mirando A, Chen K, Chen X, Kim S, Francklyn CS, Guo M. Structural basis for full-spectrum inhibition of translational functions on a tRNA synthetase. *Nat Commun*. 2015;6:6402.
- Rock FL, Mao W, Yaremchuk A, Tukalo M, Crepin T, Zhou H, et al. An antifungal agent inhibits an aminoacyl tRNA synthetase by trapping tRNA in the editing site. *Science*. 2007;316:1759–61 (80).
- Lee EY, Kim S, Kim MH. Aminoacyl-tRNA synthetases, therapeutic targets for infectious diseases. *Biochem Pharmacol*. 2018;154:424–34.
- Jain V, Sharma A. Repurposing of potent drug candidates for multiparasite targeting. *Trends Parasitol*. 2017;33(3):158–61.
- Park JK, Kim KH, Kang S, Kim W, Eom KS, Littlewood DT. A common origin of complex life cycles in parasitic flatworms: evidence from the complete mitochondrial genome of *Microcotyle sebastis* (Monogenea: Platyhelminthes). *BMC Evol Biol*. 2007;7:11.
- Zarowiecki M, Berriman M. What helminth genomes have taught us about parasite evolution. *Parasitology*. 2015;142(Suppl 1):S85–97.
- Sharma A, Sharma A. *Plasmodium falciparum* mitochondria import tRNAs along with an active phenylalanyl-tRNA synthetase. *Biochemical J*. 2015;465(3):459–69.
- Datt M, Sharma A. Evolutionary and structural annotation of disease-associated mutations in human aminoacyl-tRNA synthetases. *BMC Genomics*. 2014;15:1063.
- Bhatt TK, Kapil C, Khan S, Jairajpuri MA, Sharma V, Santoni D, Silvestrini F, Pizzi E, Sharma A. A genomic glimpse of aminoacyl-tRNA synthetases in malaria parasite *Plasmodium falciparum*. *BMC Genomics*. 2009;10:644.
- Datt M, Sharma A. Novel and unique domains in aminoacyl-tRNA synthetases from human fungal pathogens *Aspergillus Niger*, *Candida albicans* and *Cryptococcus neoformans*. *BMC Genomics*. 2014;15:1069.

30. Khan S, Sharma A, Jamwal A, Sharma V, Pole AK, Thakur KK, Sharma A. Uneven spread of cis-and trans-editing aminoacyl-tRNA synthetase domains within translational compartments of *P. falciparum*. *Scientific Rep.* 2011;1:188.
31. Kron M, Petridis M, Milev Y, Leykam J, Härtlein M. Expression, localization and alternative function of cytoplasmic asparaginyl-tRNA synthetase in *Brugia malayi*. *Mol Biochem Parasitol.* 2003;129:33–9.
32. Kron MA, Wang C, Vodanovic-Jankovic S, Zack Howard OM, Kuhn LA. Interleukin-8-like activity in a filarial asparaginyl-tRNA synthetase. *Mol Biochem Parasitol Elsevier BV.* 2012;185:66–9.
33. Ramirez BL, Howard OMZ, Dong HF, Edamatsu T, Gao P, Hartlein M, et al. *Brugia malayi* Asparaginyl-transfer RNA Synthetase induces chemotaxis of human leukocytes and activates G-protein-coupled receptors CXCR1 and CXCR2. *J Infect Dis.* 2006;193(8):1164–71.
34. Lu Z, Sessler F, Holroyd N, Hahnel S, Quack T, Berriman M, et al. Data descriptor: a gene expression atlas of adult *Schistosoma mansoni* and their gonads. *Sci Data.* 2017;4:170118.
35. Sharma A, Sharma M, Yogavel M, Sharma A. Protein translation enzyme lysyl-tRNA Synthetase presents a new target for drug development against causative agents of Loiasis and schistosomiasis. *PLoS Negl Trop Dis.* 2016;10(11):1–19.
36. Das P, Babbar P, Malhotra N, Sharma M, Jachak GR, Gonnade RG, Shanmugam D, Harlos K, Yogavel M, Sharma A, Reddy DS. Specific Stereoisomeric conformations determine the drug potency of Cladosporin scaffold against malarial parasite. *J Med Chem.* 2018;61(13):5664–78.
37. Hussain T, Yogavel M, Sharma A. Inhibition of protein synthesis and malaria parasite development by drug targeting of methionyl-tRNA synthetases. *Antimicrob Agents Chemother.* 2015;59(4):1856–67.
38. Kato N, et al. Diversity-oriented synthesis yields novel multistage antimalarial inhibitors. *Nature.* 2016;538(7625):344–9.
39. Jain V, Kikuchi H, Oshima Y, Sharma A, Yogavel M. Structural and functional analysis of the anti-malarial drug target prolyl-tRNA synthetase. *J Struct Funct Genom.* 2014;15(4):181–90.
40. Gowri VS, Ghosh I, Sharma A, Madhubala R. Unusual domain architecture of aminoacyl tRNA synthetases and their paralogs from *Leishmania major*. *BMC Genomics.* 2012;13:621.
41. Jain V, Sharma A, Singh G, Yogavel M, Sharma A. Structure-based targeting of orthologous pathogen proteins accelerates Antiparasitic drug discovery. *ACS Infect Dis.* 2017;3(4):281–92.
42. Khan S, Garg A, Camacho N, Van Rooyen J, Pole AK, Belrhali H, et al. Structural analysis of malaria-parasite lysyl-tRNA synthetase provides a platform for drug development. *Acta Crystallogr D Biol Crystallogr.* 69(5):785–95.
43. Jain V, Yogavel M, Kikuchi H, Oshima Y, Hariguchi N, Matsumoto M, et al. Targeting Prolyl-tRNA Synthetase to Accelerate Drug Discovery against Malaria, Leishmaniasis, Toxoplasmosis, Cryptosporidiosis, and Coccidiosis. *Structure.* 2017;25:1495–1505.e6 Elsevier Ltd.
44. Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite – a comprehensive resource for helminth genomics. *Mol. Biochem Parasitol.* 2017;215:2–10 Elsevier B.V.
45. Finn RD, Coghill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, et al. The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res.* 2016;44:D279–85.
46. Szymanski M, Deniziak MA, Barciszewski J. Aminoacyl-tRNA synthetases database. *Nucleic Acids Res.* 2001;29:288–90.
47. Wolf YI, Koonin EV. A tight link between orthologs and bidirectional best hits in bacterial and archaeal genomes. *Genome Biol Evol.* 2012;4:1286–94.
48. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal omega. *Mol Syst Biol.* 2011;7.
49. Talavera G, Castresana J. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol.* 2007;56:564–77.
50. Stamatakis A, Hoover P, Rougemont J. A rapid bootstrap algorithm for the RAxML web servers. *Syst Biol.* 2008;57:758–71.
51. Stamatakis A, Ludwig T, Meier H. RAxML-III: a fast program for maximum likelihood-based inference of large phylogenetic trees. *Bioinformatics.* 2005;21:456–63.
52. Huson DH, Richter DC, Rausch C, Dezulian T, Franz M, Rupp R. Dendroscope: an interactive viewer for large phylogenetic trees. *BMC Bioinformatics.* 2007;8:1–6.
53. Kumar S, Stecher G, Tamura K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* 2016;33(7):1870–4.
54. Cestari I, Stuart K. A spectrophotometric assay for quantitative measurement of aminoacyl-tRNA synthetase activity. *J Biomol Screen.* 2013;18(4):490–7.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://www.biomedcentral.com/submissions)

