

CORRECTION

Open Access



Correction: Oral exposure to bisphenol S is associated with alterations in the oviduct proteome of an ovine model, with aggravated effects in overfed females

Coline Mahé^{1*}, Marie-Emilie Lebachelier de la Riviere¹, Olivier Lasserre², Guillaume Tsikis¹, Daniel Tomas³, Valérie Labas³, Sébastien Elis¹ and Marie Saint-Dizier^{1*}

Correction: *BMC Genomics* 25, 589 (2024)
<https://doi.org/10.1186/s12864-024-10510-z>

Following publication of the original article [1] it was reported that the following text in the “Profiling of the oviduct fluid proteome according to the exposure to BPS and diet” section should be corrected:

Original text: The GO analysis of proteins in cluster 1 evidenced an enrichment in 19 biological processes (BP) and pathways, including six related to metabolism, three related to the immune system, three to the cell response to stress, and three to cell movement among the most significant ones (p-value < 0.01) (Table 1). Analysis of proteins in cluster 2 showed an enrichment in 16 BP and pathways, including two related to intracellular trafficking, five to metabolism, three to immune system, and one

related to the negative regulation of reproductive process (Table 2).

Corrected text: The GO analysis of proteins in cluster 1 evidenced an enrichment in 16 BP and pathways, including two related to intracellular trafficking, five to metabolism, three to immune system, and one related to the negative regulation of reproductive process (Table 1). Analysis of proteins in cluster 2 showed an enrichment in 18 biological processes (BP) and pathways, including six related to metabolism, two related to the immune system, three to the cell response to stress, and three to cell movement among the most significant ones (p-value < 0.01) (Table 2).

The original article has been updated.

Published online: 02 July 2024

The original article can be found online at <https://doi.org/10.1186/s12864-024-10510-z>.

*Correspondence:

Coline Mahé
coline.mahe@inrae.fr

Marie Saint-Dizier
marie.saint-dizier@inrae.fr

¹ INRAE, CNRS, Université de Tours, PRC, Nouzilly 37380, France

² INRAE, PAO, Nouzilly 37380, France

³ INRAE, Université de Tours, CHU de Tours, Plateforme de Phénotypage

Par Imagerie in/eX Vivo de L'ANIMAL À La Molécule (PIXANIM),
Nouzilly 37380, France

Reference

1. Mahé, et al. Oral exposure to bisphenol S is associated with alterations in the oviduct proteome of an ovine model, with aggravated effects in overfed females. *BMC Genomics*. 2024;25:589. <https://doi.org/10.1186/s12864-024-10510-z>.



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.