

Database

Open Access

Hmrbase: a database of hormones and their receptors

Mamoon Rashid, Deepak Singla, Arun Sharma, Manish Kumar and Gajendra PS Raghava*

Address: Bioinformatics Centre, Institute of Microbial Technology, Sector 39A, Chandigarh-160036, India

Email: Mamoon Rashid - mamoon@imtech.res.in; Deepak Singla - deep_kumar@imtech.res.in; Arun Sharma - arunsharma@imtech.res.in; Manish Kumar - manish@imtech.res.in; Gajendra PS Raghava* - raghava@imtech.res.in

* Corresponding author

Published: 9 July 2009

Received: 23 September 2008

BMC Genomics 2009, 10:307 doi:10.1186/1471-2164-10-307

Accepted: 9 July 2009

This article is available from: <http://www.biomedcentral.com/1471-2164/10/307>

© 2009 Rashid et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Hormones are signaling molecules that play vital roles in various life processes, like growth and differentiation, physiology, and reproduction. These molecules are mostly secreted by endocrine glands, and transported to target organs through the bloodstream. Deficient, or excessive, levels of hormones are associated with several diseases such as cancer, osteoporosis, diabetes etc. Thus, it is important to collect and compile information about hormones and their receptors.

Description: This manuscript describes a database called Hmrbase which has been developed for managing information about hormones and their receptors. It is a highly curated database for which information has been collected from the literature and the public databases. The current version of Hmrbase contains comprehensive information about ~2000 hormones, e.g., about their function, source organism, receptors, mature sequences, structures etc. Hmrbase also contains information about ~3000 hormone receptors, in terms of amino acid sequences, subcellular localizations, ligands, and post-translational modifications etc. One of the major features of this database is that it provides data about ~4100 hormone-receptor pairs. A number of online tools have been integrated into the database, to provide the facilities like keyword search, structure-based search, mapping of a given peptide(s) on the hormone/receptor sequence, sequence similarity search. This database also provides a number of external links to other resources/databases in order to help in the retrieving of further related information.

Conclusion: Owing to the high impact of endocrine research in the biomedical sciences, the Hmrbase could become a leading data portal for researchers. The salient features of Hmrbase are hormone-receptor pair-related information, mapping of peptide stretches on the protein sequences of hormones and receptors, Pfam domain annotations, categorical browsing options, online data submission, DrugPedia linkage etc. Hmrbase is available online for public from <http://crdd.osdd.net/raghava/hmrbase/>.

Background

According to the Medical Subject Heading (MeSH), hormones are defined as chemical substances having a specific regulatory effect on the activity of a certain organ or

organs, although the classical definition of hormones limits them to the domain of chemical signaling molecules produced by endocrine glands and secreted directly into the bloodstream. Hormones travel through the blood to

distant tissues and organs, where they can bind to specific cell sites called receptors. By binding to receptors, hormones trigger various responses in the tissues/cells containing cognate receptors [1,2]. On the basis of their chemical natures, hormones are broadly classified into protein/peptide hormones (genome-encoded) and non-peptide hormones (non-genome-encoded). Hormone-receptor interactions are amongst the most important ligand-receptor type of interactions in biological systems. The living multicellular entity depends on complex communication networks for its survival. Hormones, acting as chemical messengers, are the postmen of endocrine machinery. The endocrine system focuses on ligand-receptor interactions to play a critical role in growth and development of multicellular eukaryotes [3,4]. The data flow in this area of biological science is rapid and vast. Therefore, collection and compilation of information about these interactions, and underlying molecules (hormones and receptors), will be useful.

In recent years, efforts have been made to collect and organize receptors (like GPCRDB, ORDB, NuReBase and GRIS) [5-8]. These databases deal with different classes of receptors in biological system; for example, GPCRDB/GRIS/ORDB for G-protein coupled receptors (GPCRs) and NuReBase for nuclear hormone receptors. Various type of databases; for example SwePep [9] for endogenous peptides and PepBank [10] for peptides collected from literature using text mining tools, came into existence recently. There are a few databases which maintain information about ligands and their receptors like PRRDB [11], GLIDA [12], and EndoNet [13]. PRRDB, an immunological database, provides information regarding Pattern Recognition Receptors and their ligands. GLIDA is developed with possible implications in chemical genomic research and GPCR-related drug discovery, whereas EndoNet is an information resource about intercellular regulatory communication. Though existing databases provide important information, there is lack of a comprehensive resource on hormones and their receptors.

In order to complement existing databases in the field, and to understand hormones and their interaction with receptors, we have developed a database called Hmrbase. This database provides comprehensive information about hormones and receptors. Various data fields like hormone precursor, subcellular localization, post-translational modification, taxonomy, source organism, function, description, tissue specificity, molecular weight, similarity to other proteins, and mapping of hormone peptide on its corresponding precursor etc. have been included for peptide hormones and their receptor. For non-peptide hormones, the data fields consist of their names, molecular weights and molecular formulae, IUPAC names, canonical and isomeric smile formulae, melting points, LogP val-

ues, water solubility, and their corresponding receptors etc. Various co-ordinate files such as PDB, SDF, and MOL files are available for download. Structure visualization tools such as Advance Chemistry Development (ACD) structure drawing applet [14] (for 2-D visualization) and Jmol applet [15] (for 3-D visualization) have been embedded in Hmrbase. Links to neighbors (external links) like Swiss-Prot [16], PDB [17], NCBI Gene Database [18], Pfam [19], PubChem [20], KEGG [21], HMDB [22], DrugBank [23], and DrugPedia [24] have been incorporated in Hmrbase to make it a complete system. Moreover, hormones and receptors entries are linked to their corresponding receptors and hormones, respectively. Sequence similarity search, peptide mapping and domain search tool, in case of protein hormone and receptor, facilitates the extraction of useful information. In addition to text search, a structural similarity-based search option for non-peptide hormones supports the search algorithm. Thus, Hmrbase provides both comprehensive and easy-to-use information related to hormones and their receptors.

Construction and content

For collection of peptide hormones, we have extensively searched Swiss-Prot, other databases and the related literature. Initially, we started searching through Sequence Retrieval System (SRS) of Swiss-Prot with a keyword "hormone" against the "Description" field with wild card. Then we exploited GPCRDB for hormone receptor. GPCR class A comprises hormone receptors such as serotonin, cholecystokinin, melanocortin, prolactin, somatostatin, vasopressin, adrenomedullin, melanin etc. GPCR class B comprises calcitonin, glucagons, diuretic, parathyroid, secretin hormone receptors. Regarding collection of non-peptide hormones we searched various databases like PubChem, Human Metabolome Database (HMDB), and EndoNet. The corresponding receptors were taken from PubChem, literature databases like PubMed, EndoNet, DrugBank, NuclearDB [25], and Swiss-Prot. Detailed information about data collection and manual curation has been included in the additional file 1.

This database consists of extensively manually curated information about peptide hormones, non-peptide hormones, and their receptors. The statistics of the data has been given in Table 1 and Table 2. Following is the brief description of data available at Hmrbase.

Peptide hormones (PH)

Different PHs from various taxonomical classes have been collected and compiled. At present the database encapsulates 1585 PHs. A pool of information has been supplied with each hormone entry. The type of information has been explained under "data structure" heading.

Table 1: Hmrbase data types and their corresponding numbers

Data types		Number of entries	Total
Hormones	Peptide/protein	1585	1955
	Non-peptide	370	
Receptors (for)	Peptide hormone	828	2996
	Non-peptide hormone	2168	
Hormone-Receptor Pair	Peptide hormone-Receptor	569	4121
	Non-peptide hormone-Receptor	3552	

Non-Peptide hormones (NPH)

These are basically small chemicals and play an important role in signal transduction pathways regulating complex networks of gene expression. A total of 370 such molecules have been compiled in Hmrbase.

Receptors for peptide hormones (RPH)

Altogether, there are 828 receptor entries for peptide hormones. Mostly, these are G-protein coupled receptors

(GPCR) on the membranes of cell surfaces, which sense external stimuli (in the form of ligands) to transduce the information to intracellular region.

Receptors for non-peptide hormones (RNH)

Receptors for non-peptide hormones are mainly ligand-activated nuclear transcription factors. These are actively involved in alterations of gene expression which, in turn,

Table 2: Some of the biological taxa with corresponding number of entries in Hmrbase database

Important taxa	Number of entries in database	
	Hormone (%)	Receptor (%)
Chordate	1362 (69.6)	2309 (77.06)
Arthropoda	198 (10.1)	207 (6.90)
Mollusca	19 (0.97)	33 (1.10)
Metatheria	29 (1.48)	15 (0.50)
Eutheria	753 (38.51)	1391 (46.42)
Primates	170 (8.69)	478 (15.95)
Rodents	227 (11.61)	535 (17.85)
Cetartiodactyla (whales, hippos, ruminants, pigs, camels)	214 (10.94)	239 (7.97)
New world monkeys	13 (0.66)	49 (1.63)
Old world monkeys	44 (2.25)	69 (2.30)
Gorilla	5 (0.25)	3 (0.10)
Chimpanzee	14 (0.71)	14 (0.46)
Human	84 (4.29)	310 (10.34)

Values in parentheses are the percentage of hormones or receptors for a particular taxon included in Hmrbase database

regulate the normal physiology of an organism. A total of 2168 RNH have been maintained in Hmrbase.

Hormone-receptor pair

To understand the functional diversity and the mode of action of any hormone, information about its receptor is essential. Approximately, 4121 hormone-receptor functional interactions have been incorporated in Hmrbase.

Utility

Web Tools

Apart from the collection of hormone molecules and their receptors, a wide variety of information can be generated using the online software/tools provided with Hmrbase database. Following are the main web tools provided with the Hmrbase database:

Keyword Search

There are separate search pages for hormones and receptors. Both are almost identical in architecture except for few data fields. Searching can be performed on any field separately, or on all fields simultaneously, using a specific keyword, to retrieve data from the database. The search option is restricted to almost all the data fields available

in the database. User can define which search results are to be displayed.

Structural Similarity Search

Hmrbase also provides a tool to search for entries related to non-peptide hormones, based on their structural similarities. JAVA Molecular Editor (JME) [26] along with JC search tool [27] facilitates the structure based searching of the Hmrbase entries. Structures drawn in the JME editor are used by JC search tool to search for similar structures in the database. Users can also select one of the different search types e.g. substructure search, exact search, superstructure search etc. Figure 1 illustrates an example of structure based search of Hmrbase entries where a phenyl group was used as a query substructure. Structure search resulted in two entries namely 16, 17-epiestriol and 6-keto estriol hormones.

2-D and 3-D structure visualization tools

For non-peptide hormones, Hmrbase is facilitated by structure display tools. Jmol and Advanced Chemistry Development (ACD) structure drawing applets have been implemented, to show 3-D and 2-D structures, respectively. Figure 2 illustrates a 3-D structure of testosterone




Figure 1

An example of structural similarity based search of Hmrbase entries. An example of structure based search of Hmrbase entries where a phenyl group was used as a query substructure. Structure search resulted in two entries namely 16, 17-epiestriol and 6-keto estriol hormones.

hormone displayed using the Jmol applet embedded in Hmrbase.

DrugPedia

Hmrbase provides a DrugPedia link corresponding to each entry. Data stored in the Hmrbase platform follow a definite pattern. Therefore, to provide more flexibility to different kinds of data, a DrugPedia link would be beneficial in which users can update or add any relevant information. The important and relevant information from the DrugPedia page may be included in the main database frame after validation during the updation of Hmrbase. Figure 3 shows a typical DrugPedia page for testosterone hormone.

Sequence Similarity Search

A customized BLAST [28] tool has been made available which searches a user-defined query against the sequence of the hormone, or receptor, or both. It may be useful in characterization of orphan receptors and fishing out of homologous sequences from the database, based on sequence similarity.

Peptide Mapping

Users can map active subsequences or stretches of amino acids on the hormone and/or receptor protein sequences. This will add to the information regarding the distribu-

tion of functional stretches of amino acid over the entire hormone and/or the receptor sequences present in the database. Such a type of mapping might be useful in understanding the functional diversities of biologically active peptides. Peptide mapping in Hmrbase is simply implemented using "exact string search", which searches user defined queries for peptides in hormone/receptor protein sequences.

Pfam Domain Search

A group of functionally similar sequences may represent common peptide domain(s). This concept underlies the domain-wise classifications of protein families and sub-families. Therefore, the types of domain(s) present in a hormone, and in receptor protein sequences, determine their classification. Thus, the Pfam domain search algorithm has been embedded into the Hmrbase platform. It has been observed that some domains are highly frequent and some are rare amongst Hmrbase entries. All the unique domains from peptide hormone and receptor entries along with their frequency distribution have been shown in Table 3.

Data Structure

Several tables in relational fashion facilitate the data architecture of Hmrbase. There are three primary tables at the front, one each for peptide hormones, non-peptide




Figure 2

Structure visualization by Jmol. Structure of testosterone hormone as visualized in Jmol, showing different labels of atoms; white balls are hydrogen atom, red oxygen, and gray are carbon atoms.

Testosterone

Testosterone is a **steroid hormone** from the **androgen** group. In mammals, testosterone is primarily secreted in the **testes** of males and the **ovaries** of females, although small amounts are also secreted by the **adrenal glands**. It is the principal **male sex hormone** and an **anabolic steroid**. Its production is stimulated by **LUTEINIZING HORMONE** from the **PITUITARY GLAND**. In turn, testosterone exerts feedback control of the pituitary LH and FSH secretion. Depending on the tissues, testosterone can be further converted to **DIHYDROTESTOSTERONE** or **ESTRADIOL**.

In both men and women, testosterone plays a key role in health and well-being as well as in sexual functioning. Examples include enhanced **libido**, increased energy, increased production of red blood cells and protection against **osteoporosis**. On average, an adult **human male** body produces about forty to sixty times more testosterone than an adult female body, but females are, from a behavioral perspective (rather than from an anatomical or biological perspective), more sensitive to the hormone. <ref>James McBride Dabbs, 2000</ref> However the overall ranges for male and female are very wide, such that the ranges actually overlap at the low end and high end respectively.

Contents [hide]

- 1 History
- 2 Physicochemical Properties
 - 2.1 Melting Point
 - 2.2 log P
 - 2.3 Water Solubility
 - 2.4 Vapor Pressure
 - 2.5 Henry's Law Constant
 - 2.6 Atmospheric OH Rate Constant
- 3 Production
 - 3.1 Natural
 - 3.2 Artificial
 - 3.2.1 Esterification
- 4 Virilizing and effects on humans
 - 4.1 Prenatal androgen effects
 - 4.2 Early infancy androgen effects
 - 4.3 Early postnatal effects
 - 4.4 Advanced postnatal effects
 - 4.5 Adult testosterone effects
- 5 Effects on the brain
- 6 Mechanism
- 7 Therapeutic use
 - 7.1 Routes of administration

Chemical data

Formula	C ₁₉ H ₂₈ O ₂
Mol. mass	288.42
SMILES	eMolecules & PubChem
Physical data	
Melt. point	155°deg C EXP °C
Solubility in water	23.4 mg/L at 25°C EXP mg/mL
Spec. rot	+110.2°
SEC Combust	-11080 kJ/mol
Pharmacokinetic data	
Bioavailability	low (due to extensive first pass metabolism)
Metabolism	Liver, Testis and Prostate
Half life	2-4 hours
Excretion	Urine (90%), feces (6%)
Therapeutic considerations	
Pregnancy cat.	X (USA), Teratogenic effects

Figure 3

A typical DrugPedia page for testosterone hormone. Figure shows a typical DrugPedia page for testosterone hormone. Top right corner contains link for new account creation and login. To edit, discuss, or see history of a particular page, links are embedded on top left of the page.

hormones and receptors. The query from the user is replied to with the help of these three tables. The table for peptide hormones provides a 'series' of information like hormone name, active sequence, length of active sequence, mapping of active sequence on its precursor, related references, function, Swiss-Prot accession number, precursor/protein description, source organism, taxonomy, subcellular localization, developmental stage, similarity, tissue specificity, post-translational modification, length and molecular weight (in Dalton) of precursor/source. Each non-peptide hormone entry presents the name, description, IUPAC name, canonical and isomeric smile, melting point, LogP, water solubility, molecular weight, molecular formula, corresponding receptor, and external links to different databases such as PubChem, PDB, KEGG, HMDB, DrugBank etc. Each entry about a hormone is internally linked to its corresponding receptor entry in the receptor table. Moreover, each of the Hmrbase entries has been linked to DrugPedia (a Wikipedia for Drug Discovery), where users may contribute data and updated information. Further, FASTA sequences can be downloaded by clicking the link supplied by hormone and receptor tables. The table for receptors supplies more

or less the same information as peptide hormone table. Bi-directional cross-linking has been provided between hormone entry and its corresponding receptor entry. Figure 4 illustrates the whole data structure of Hmrbase.

Data Flow

There are separate search pages to query hormone, or receptor, entries. Keyword search options operate on a variety of data types like hormone name, description, developmental stage, source organism, functions etc. Each hormone entry is linked to its corresponding receptors (if available). In a similar way the receptor search result page leads towards their corresponding ligand(s). This scheme of data flow is depicted in Figure 5. The keyword "ghrelin" fetched a total of 20 entries (Figure 5, top left panel) from hormone table of Hmrbase. One of the entries out of 20 was for Swiss-Prot accession number Q9GKY5, showing corresponding receptor entry (P34999) along with other data fields (Figure 5, top right panel). Figure 5 (bottom panel) showed complete page for P34999 from receptor table. The Hmrbase entries provide links to many external databases such as Swiss-Prot, NCBI Gene, PDB, Pfam etc. Moreover, each and every entry was also linked to Drug-

Table 3: Unique Pfam domains and their occurrence among Hmrbase entries

Serial no.	Peptide Hormone		Receptor	
	Pfam Domain Name	Frequency	Pfam Domain Name	Frequency
1.	Insulin	253	zf-C4	1593
2.	Hormone_1	170	Hormone_recep	1459
3.	ACTH_domain	166	7tm_1	824
4.	Op_neuropeptide	156	7tm_2	129
5.	Hormone_2	144	HRM	119
6.	Gastrin	96	LRR_1	42
7.	Cys_knot	78	fn3	34
8.	Somatostatin	55	DUF1856	13
9.	Hormone_6	54	EGF	7
10.	Thymosin	46	EGF_CA	7
11.	Crust_neurohorm	45	Laminin_G_2	6
12.	Hormone_3	33	Laminin_G_1	6
13.	CRF	24	Furin-like	4
14.	Motilin_assoc	20	Recep_L_domain	4
15.	Motilin_ghrelin	20	Ldl_recept_a	4
16.	ELH	18	Calsequestrin	1
17.	Bombesin	11	Homeobox	1
18.	TRH	10	Thioredoxin	1
19.	Hormone_5	9		
20.	Adipokin_hormo	8		
21.	Eclosion	3		
22.	Crust_neuro_H	2		

Frequency of a domain is the number of entries in Hmrbase database harboring that domain.

Pedia, where one can contribute recent or updated information.


Web Interface and Application

Simple HTML and CSS technologies have been used to build the static web interface. MySQL, an object-relational Database Management System (RDBMS), works at the backend. Server-side scripting makes use of PHP. The whole software system runs on IBM SAS x3800 machine

under RedHat Enterprise Linux 5 environment using Apache httpd server. PHP and MySQL combination is quite efficient and powerful for database management.

Discussion and conclusion

Hmrbase is a comprehensive information resource about hormones and their receptors. Ligand-receptor interactions have been elucidated in a bi-directional manner. Users can start from searching hormone entry (ies) and

**Figure 4**

Data structure of Hmrbase. Data types for hormone and receptor tables have been shown in different color schemes; black colored data types/tools are common for hormones as well as receptors, brown colored are exclusively for hormones.

end up in their corresponding receptor (s); and vice-versa. The database will fulfill the requirements of theoretical as well as clinical endocrinologists. Data structure of Hmrbase is quite simple and convenient for general users. The mature hormone sequence is mapped on its precursor protein sequence in order to define the functional modes of hormones. Furthermore, this information can be exploited by experimental scientists to design better ligands for a particular receptor or for studying binding affinity of hormones to their corresponding receptors. Protein domains have been inferred as the basic building blocks of protein interactions [29]. Therefore, to derive the Pfam domains distribution among the entries of Hmrbase, a domain search facility has been embedded.

Furthermore, a collection of non-peptide hormone molecules along with various operational tools such as ACD/Structure drawing applet, Jmol, and JC Search with JME editor facilitates the completeness of the database.

Hopefully this customized database will expand quantitatively as well as qualitatively in coming days to cover the annotation gaps such as orphan receptors and probably any novel hormone molecule.

Application of Hmrbase

Hmrbase presents data in a sophisticated way. Apart from text search, several browsing options facilitate the retrieval of important datasets, such as entries for a particular organism or hormone name or specific domain or domain combinations etc. Moreover, hormone-receptor and receptor-hormone pairs have been presented to infer the range of action of a particular hormone or receptor. Different types of structure search algorithm such as substructure, exact, superstructure search would help in compiling the set of entries containing a particular functional group or moiety. Moreover, each entry of Hmrbase has been linked to DrugPedia, which would serve as a complement to Hmrbase entries. Any new or updated information posted on DrugPedia would be included in Hmrbase

HMRbase: A database of hormones

Home | Hormone | Receptor | Tools

Keyword: ghrelin
Search Type: Similar
Database: Peptide Hormone

Go Page No. 2

Go Query Page Result No - 7 of 20

HMRbase Accession No.	10115
Swiss-prot Accession No.	Q9GKY5 (Sequence in FASTA format)
Description	Appetite-regulating hormone precursor (Growth hormone secretagogue)(Growth hormone-releasing peptide)(Motilin-related peptide)(Contains: Ghrelin; Obestatin).
Origin	Sus scrofa (Pig).
Taxonomy	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Laurasiatheria; Cetartiodactyla; Suina; Suidae; Sus.
Sub-location	Secreted protein (By similarity).
Developmental Stage	N/A
Similarity	Belongs to the motilin family.
Tissue Specificity	N/A
Post translational modification	O-6-octanoylation is essential for ghrelin activity (By similarity). Amidation of Leu-99 is essential for obestatin activity (By similarity).
Function	Ghrelin is the ligand for growth hormone secretagogue receptor type 1 (GHSR). Induces the release of growth hormone from the pituitary. Has an appetite-stimulating effect, induces adiposity and stimulates gastric acid secretion. Involved in growth regulation
Protein Length	118 Amino acids
Molecular weight	12786
Reference	1 Kojima M.; Submitted (DEC-1999) to the EMBL/GenBank/DBJ databases. 2 Rossetti J., Lacroix D., Dubreuil P.; Submitted (MAR-2001) to the EMBL/GenBank/DBJ databases. 3 Ying M., Yang Z.; Submitted (JAN-2006) to the EMBL/GenBank/DBJ databases.
Domain Name	Motilin, assoc. Motilin, ghrelin
Hormone Name	Ghrelin
Hormone Sequence	GSSFLSPHQKVVQRKESKKPAALKPR
Hormone Map	28 Residues from position (25-52)
Receptor	P34999
Gene ID	296728
PDB ID	

HMRbase: A database of hormones and their receptors

Home | Hormone | Receptor | Tools | Help | Contact

Receptor Database

HMRbase Accession No.	21035
Swiss-prot Accession No.	P34999 (Sequence in FASTA format)
Description	Growth hormone-releasing hormone receptor precursor (GHRH receptor)(GRF receptor) (GRFR).
Source Organism	Sus scrofa (Pig).
Taxonomic Classification	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Laurasiatheria; Cetartiodactyla; Suina; Suidae; Sus.
Subcellular Location	Membrane; multi-pass membrane protein.
Developmental Stage	N/A
Similarity	Belongs to the G-protein coupled receptor 2 family.
Tissue Specificity	Pituitary gland. Also detected in the lymphocytes and thymocytes.
Post translational modification	N/A
Function	Receptor for GRF, coupled to G proteins which activate adenyl cyclase. Stimulates somatotroph cell growth, growth hormone gene transcription and growth hormone secretion.
Protein Length	423 Amino acids
Molecular weight	47200
Domain Name	7tm_2_HRM
Reference	1 PubMed abstract 8413847 2 Hazem H.A., Zhang X., Smith D.P., Heiman M.L., Hsiung H.M.; Submitted (APR-1996) to the EMBL/GenBank/DBJ databases.
Hormone	P01287 Q9GKY5
Gene ID	397100
PDB ID	

Figure 5

Data flow in Hmrbase. Hmrbase hormone search result page using keyword "ghrelin" as description, a total of 20 entries were found (top left panel); out of 20 hormone entries, one complete entry has been shown for Swiss-Prot accession number Q9GKY5 and its receptor counterpart is P34999 (top right panel); by clicking the link for P34999 a complete entry page for receptor P34999 appears and among its ligand counterparts is Q9GKY5 (bottom panel).

database after validation. Thus Hmrbase would be a comprehensive and stable system for biomedical researchers and bioinformatician.

Limitations and future prospects

Several new data types such as pharmacological data are being collected to incorporate into Hmrbase. The major limitation of this resource is the lack of a fully automated database populating system. Nevertheless, we have devised different models for updation of Hmrbase at a period of every three month (see Additional file 1).

Availability and requirements

Hmrbase is available at: <http://crdd.osdd.net/raghava/hmrbase/>. To access all features of Hmrbase database to its

optimum level, javascript and Java Runtime Environment (JRE) plugin must be enabled.

Authors' contributions

MR collected and compiled the data from literature and public databases. MR made the structure of the database. DS collected non-peptide hormone molecules and their nuclear receptor counterparts. MK and MR incorporated various amino acid sequence manipulative tools in Hmrbase. Web interface was designed by AS and MR. AS contributed in converting Perl scripts into PHP and PostgreSQL tables into MySQL tables originally made by MR and MK. AS and DS incorporated tools for non-peptide hormone manipulation. GPSR conceived the project, coordinated it and refined the manuscript drafted by MR.

Additional material

Additional file 1

Supplementary material. It consists of detail strategy of data collection in Hmrbase and the updation schemes.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-10-307-S1.pdf>]

Acknowledgements

Authors are grateful to Dr P. Guptasarma for critically reading the manuscript. We are also thankful to Mr. Hifzur Rahman Ansari and Mr. Harinder for their help in preparing manuscript. The authors are thankful to Council of Scientific and Industrial Research (CSIR) and Department of Biotechnology, Government of India, for financial assistance. This report has IMTECH communication number 12/2008.

References

- Lodish HBA, Zipursky SL, Matsudaira P, Baltimore D, Darnell J: **Molecular Cell Biology**. 4th edition. NY: W.H. Freeman and Company; 2000.
- Nussey SSWS: **Endocrinology: An Integral Approach**. Oxford: BIOS Scientific Publishers Ltd; 2001.
- Jones DS, Silverman AP, Cochran JR: **Developing therapeutic proteins by engineering ligand-receptor interactions**. *Trends in Biotechnology* 2008, **26(9)**:498-505.
- Łącka K, Czyżyk A: **Hormones and the cardiovascular system**. *Endokrynol Pol* 2008, **59(5)**:420-433.
- Horn F, Bettler E, Oliveira L, Campagne F, Cohen FE, Vriend G: **GPCRDB information system for G protein-coupled receptors**. *Nucleic Acids Res* 2003, **31(1)**:294-297.
- Crasto C, Marengo L, Miller P, Shepherd G: **Olfactory Receptor Database: a metadata-driven automated population from sources of gene and protein sequences**. *Nucleic Acids Res* 2002, **30(1)**:354-360.
- Ruau D, Duarte J, Ourjda T, Perriere G, Laudet V, Robinson-Rechavi M: **Update of NUREBASE: nuclear hormone receptor functional genomics**. *Nucleic Acids Res* 2004:D165-167.
- Van Durme J, Horn F, Costagliola S, Vriend G, Vassart G: **GRIS: glycoprotein-hormone receptor information system**. *Mol Endocrinol* 2006, **20(9)**:2247-2255.
- Falsh M, Skold K, Norrman M, Svensson M, Fenyo D, Andren PE: **SwePep, a database designed for endogenous peptides and mass spectrometry**. *Mol Cell Proteomics* 2006, **5(6)**:998-1005.
- Shtatland T, Guettler D, Kossodo M, Pivovarov M, Weissleder R: **PepBank – a database of peptides based on sequence text mining and public peptide data sources**. *BMC Bioinformatics* 2007, **8**:280.
- Lata S, Raghava GP: **PRRDB: a comprehensive database of pattern-recognition receptors and their ligands**. *BMC Genomics* 2008, **9**:180.
- Okuno Y, Yang J, Taneishi K, Yabuuchi H, Tsujimoto G: **GLIDA: GPCR-ligand database for chemical genomic drug discovery**. *Nucleic Acids Res* 2006:D673-677.
- Potapov A, Lieblich I, Donitz J, Schwarzer K, Sasse N, Schoeps T, Crass T, Wingender E: **EndoNet: an information resource about endocrine networks**. *Nucleic Acids Res* 2006:D540-545.
- Advanced Chemistry Development (ACD) Structure Drawing Applet (SDA) [<http://www.acdlabs.com/products/java/sda/>]
- Jmol: an open-source Java viewer for chemical structures in 3D [<http://www.jmol.org/>]
- Boeckmann B, Blatter MC, Famiglietti L, Hinz U, Lane L, Roehert B, Bairoch A: **Protein variety and functional diversity: Swiss-Prot annotation in its biological context**. *C R Biol* 2005, **328(10-11)**:882-899.
- Zardecki C: **Interesting structures: education and outreach at the RCSB Protein Data Bank**. *PLoS Biol* 2008, **6(5)**:e117.
- Maglott D, Ostell J, Pruitt KD, Tatusova T: **Entrez Gene: gene-centered information at NCBI**. *Nucleic Acids Res* 2005:D54-58.
- Finn RD, Tate J, Mistry J, Coghill PC, Sammut SJ, Hotz HR, Ceric G, Forslund K, Eddy SR, Sonnhammer EL, et al.: **The Pfam protein families database**. *Nucleic Acids Res* 2008:D281-288.
- Han L, Wang Y, Bryant SH: **Developing and validating predictive decision tree models from mining chemical structural fingerprints and high-throughput screening data in PubChem**. *BMC Bioinformatics* 2008, **9**:401.
- Kanehisa M, Araki M, Goto S, Hattori M, Hirakawa M, Itoh M, Katayama T, Kawashima S, Okuda S, Tokimatsu T, et al.: **KEGG for linking genomes to life and the environment**. *Nucleic Acids Res* 2008:D480-484.
- Wishart DS, Tzur D, Knox C, Eisner R, Guo AC, Young N, Cheng D, Jewell K, Arndt D, Sawhney S, et al.: **HMDB: the Human Metabolome Database**. *Nucleic Acids Res* 2007:D521-526.
- Wishart DS, Knox C, Guo AC, Cheng D, Shrivastava S, Tzur D, Gautam B, Hassanali M: **DrugBank: a knowledgebase for drugs, drug actions and drug targets**. *Nucleic Acids Res* 2008:D901-906.
- DrugPedia: A Wikipedia for Drug Discovery [http://crdd.osdd.net/drugpedia/index.php/Main_Page]
- Horn F, Vriend G, Cohen FE: **Collecting and harvesting biological data: the GPCRDB and NucleaRDB information systems**. *Nucleic Acids Res* 2001, **29(1)**:346-349.
- JAVA MOLECULAR EDITOR [<http://www.molinspiration.com/jme/>]
- Csizmadia F: **JChem: Java Applets and Modules Supporting Chemical Database Handling from Web Browsers**. *Journal of Chemical Information and Computer Sciences* 2000, **40(2)**:323-324.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: **Gapped BLAST and PSI-BLAST: a new generation of protein database search programs**. *Nucleic Acids Res* 1997, **25(17)**:3389-3402.
- Deng M, Mehta S, Sun F, Chen T: **Inferring domain-domain interactions from protein-protein interactions**. *Genome Res* 2002, **12(10)**:1540-1548.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

