

RESEARCH ARTICLE

Open Access

Analysis of the heat shock response in mouse liver reveals transcriptional dependence on the nuclear receptor peroxisome proliferator-activated receptor α (PPAR α)

Beena Vallanat¹, Steven P Anderson², Holly M Brown-Borg³, Hongzu Ren¹, Sander Kersten⁴, Sudhakar Jonnalagadda⁵, Rajagopalan Srinivasan^{5,6}, J Christopher Corton^{1*}

Abstract

Background: The nuclear receptor peroxisome proliferator-activated receptor alpha (PPAR α) regulates responses to chemical or physical stress in part by altering expression of genes involved in proteome maintenance. Many of these genes are also transcriptionally regulated by heat shock (HS) through activation by HS factor-1 (HSF1). We hypothesized that there are interactions on a genetic level between PPAR α and the HS response mediated by HSF1.

Results: Wild-type and PPAR α -null mice were exposed to HS, the PPAR α agonist WY-14,643 (WY), or both; gene and protein expression was examined in the livers of the mice 4 or 24 hrs after HS. Gene expression profiling identified a number of *Hsp* family members that were altered similarly in both mouse strains. However, most of the targets of HS did not overlap between strains. A subset of genes was shown by microarray and RT-PCR to be regulated by HS in a PPAR α -dependent manner. HS also down-regulated a large set of mitochondrial genes specifically in PPAR α -null mice that are known targets of PPAR γ co-activator-1 (PGC-1) family members. Pretreatment of PPAR α -null mice with WY increased expression of PGC-1 β and target genes and prevented the down-regulation of the mitochondrial genes by HS. A comparison of HS genes regulated in our dataset with those identified in wild-type and HSF1-null mouse embryonic fibroblasts indicated that although many HS genes are regulated independently of both PPAR α and HSF1, a number require both factors for HS responsiveness.

Conclusions: These findings demonstrate that the PPAR α genotype has a dramatic effect on the transcriptional targets of HS and support an expanded role for PPAR α in the regulation of proteome maintenance genes after exposure to diverse forms of environmental stress including HS.

Background

Physiological and chemical stresses produce disease states in which proteins are damaged or misfolded in part through increases in oxidative stress. Many endogenous pathways are engaged in restoring cellular homeostasis, including stabilization of unfolded proteins to prevent aggregation and removing damaged or excess proteins through proteolysis. Stabilization of unfolded proteins is performed by molecular chaperones that

assist in the folding of nascent polypeptides. Many genes encoding for chaperones exhibit increased expression after exposure to a wide variety of stimuli including chemical exposure or increased temperatures and are thus called heat shock (HS) protein (*Hsp*) genes [1-3]. These proteins play key roles in a number of human diseases [4]. Expression of some *Hsp* is essential for cellular survival under physical or chemical exposure conditions that increase oxidative stress [5,6]. Regulation of the *Hsp* genes by heat or chemical-induced oxidative stress is controlled in part by HS factor 1 (HSF1), activated under conditions in which the level of unfolded proteins increase [1-3]. Microarray studies of mouse

* Correspondence: corton.chris@epa.gov

¹NHEERL Toxicogenomics Core, US EPA, Research Triangle Park, NC 27711, USA

embryonic fibroblasts from wild-type and HSF1-null mice or a human cervical carcinoma cell line have shown that HSF1 controls only a subset of the genes altered by HS [7,8] indicating that other inducible pathways play roles in regulating the *Hsp* genes.

The nuclear receptor peroxisome proliferator-activated receptor α (PPAR α) is one of three PPAR subtypes that regulate lipid and glucose homeostasis, tissue growth and inflammation after exposure to a large class of structurally heterogeneous industrial and pharmaceutical chemicals called peroxisome proliferator chemicals (PPC) [9,10]. The PPAR α subtype plays a key role in mediating the effects of hypolipidemic and xenobiotic PPC in liver, kidney, heart and skin. Exposure to PPC leads to regulation of a large number of genes including up-regulation of those involved in lipid homeostasis and down-regulation of inflammatory genes that are generally abolished in PPC-exposed PPAR α -null mice [11-13].

There is compelling evidence that PPAR α protects tissues from chemical-induced oxidative stress (reviewed in [14]). Prior exposure of rats and mice to PPC protects the liver from damage by cytotoxic agents that induce oxidative stress [15]. The hypolipidemic drug and PPC, clofibrate, protects the liver from damage by the cytotoxicant acetaminophen in wild-type but not PPAR α -null mice [16]. Compared to wild-type mice, untreated PPAR α -null mice or primary hepatocytes isolated from PPAR α -null mice were more sensitive to carbon tetrachloride-, paraquat- or cadmium-induced toxicity [12]. The beneficial effects of caloric restriction in protecting the liver from cytotoxicant-induced liver injury were shown to depend on PPAR α [17]. Additional studies have shown that PPAR α plays a positive role in recovering from partial hepatectomy [18]. In the kidney, PPAR α -null mice were more sensitive to damage after ischemia-reperfusion injury, and prior exposure of wild-type mice to PPC reduces the injury [19,20]. Our previous microarray studies identified an overlap in the genes regulated by the PPAR α agonist WY-14,643 (WY) and those regulated by HS through HSF1 [12]. The proteome maintenance genes included those involved in protein folding (e.g., *Hsp* genes) as well as ubiquitin-dependent and -independent proteolytic processing through the proteasome (e.g., *Psm* genes). Altered regulation of these genes by PPC could help to explain why PPC exposure through PPAR α helps to protect tissues from environmental stressors.

In the present study, we hypothesized that there were interactions on a genetic level between PPAR α and the HS response mediated by HSF1. We dissected the contribution of PPAR α by examining gene expression changes in the livers of wild-type and PPAR α -null mice after HS. We found that a number of genes regulated by HS were dependent on PPAR α for regulation.

Furthermore, we showed that PPAR α -null mice were more sensitive to the transcriptional effects of HS and exhibited a remarkably different transcriptional response compared to wild-type mice. Our findings suggest PPAR α is a major regulator of stress responses in the liver.

Methods

Animals, exposure to heat and WY

All animal studies were carried out at CIIT Centers for Health Research, Research Triangle Park, NC. Studies utilized wild-type and PPAR α -null male mice 9-12 weeks of age on a mixed SV129/C57BL/6 background. These mice have been described previously [21]. The mice were originally obtained from Dr. Frank Gonzalez, NCI, National Institutes of Health to establish a breeding colony at CIIT. Control and treated mice were provided with NIH-07 rodent chow (Zeigler Brothers, Gardeners, PA) and deionized, filtered water *ad libitum*. Lighting was on a 12-hr light/dark cycle. Mice (n = 5-6 per group) were fed a control diet or a diet containing the PPAR α agonist WY-14,643 (ChemSyn Science, Lenexa, KS) (WY) in the diet (500 ppm) for 1 week followed by a 40-min heat stress at 42°C or held at room temperature in wire rack cages. Mice were sacrificed 4 hrs, 24 hrs and 3 days after the heat stress.

Heat in the exposure chamber was generated by a combination of the building reheat system, an in-line air heater and two small surface heaters. The in-line heater (Riley Equipment Company, Inc., Houston, TX) was set to approximately 300°F. The heated air traveled through a 2 foot by 2 inch inlet tube constructed of PVC, stainless steel and Teflon into the bottom of the H1000 inhalation chamber. A small diverter was placed in the bottom of the H1000 inhalation chamber to prevent a chimney effect in the chamber and increasing the efficiency of the heat distribution. The stainless steel animal cage unit was placed in the center of the H1000 inhalation chamber and a feces/urine catch pan was placed in the chamber. A small (1500 watt) heater was placed on either side of the H1000 inhalation chamber approximately 4 inches from the surface. These heaters heated the stainless steel sides and heated the surface enough to allow the internal temperature of the chamber to reach approximately 42°C. The temperature was monitored in three locations of the H1000 inhalation chamber: front door above the animal cage unit, back door below the animal cage unit and in the center using temperature probes (Rotronic Hygrometer Series 1200, Rotronic Instrument Corp., Huntington, NY).

Portions of the livers were rapidly snap-frozen in liquid nitrogen and stored at -70°C until analysis. Where appropriate, slices of liver were fixed in 10% neutral buffered formalin for 48 h, transferred to 70% ethanol, and

embedded in paraffin; 5- μ m sections were cut and mounted on slides and stained with H&E. H&E-stained liver sections were examined by light microscopy. Clinical chemistry markers in the serum from animals were measured by Antech Diagnostics, Charlotte, NC. All animal studies were conducted under federal guidelines for the use and care of laboratory animals and were approved by Institutional Animal Care and Use Committees.

RNA Isolation and Analysis of Gene Expression

Liver RNA was isolated using a modified guanidium isothiocyanate method (TRIzol[®]; Invitrogen) and was further purified using silica membrane spin columns (RNeasy[®]; Qiagen, Valencia, CA). RNA integrity was assessed by the RNA 6000 LabChip[®] kit using a 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA). Global gene expression changes were examined using the Affymetrix platform. Gene expression changes were assessed in the livers from three mice in each of the 8 treatment groups in wild-type and PPAR α -null mice using a total of 24 chips. Gene expression in each animal was assayed on a separate chip. Biotin-labeled cRNA was produced from 15 μ g total RNA using an Affymetrix “one-way” labeling kit. Total cRNA was quantified using a Nano-Drop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE) and evaluated for quality after fragmentation on a 2100 Bioanalyzer. Following overnight hybridization at 45°C to Affymetrix U74Av2 GeneChips in an Affymetrix Model 640 GeneChip hybridization oven, the arrays were washed and stained using an Affymetrix 450 fluidics station as recommended by the manufacturer and scanned on an Affymetrix Model 3000 scanner. After scanning, raw data (Affymetrix .cel files) were obtained using Affymetrix GeneChip Operating Software (version 1.4). All of the Affymetrix (Santa Clara, CA) .cel files were analyzed by Bioconductor SimpleAffy to assess data quality [22].

Data (.cel files) was analyzed and statistically filtered using Rosetta Resolver[®] version 7.1 software (Rosetta Inpharmatics, Kirkland, WA). Statistically significant genes were identified using one-way ANOVA with a false discovery rate (Benjamini-Hochberg test) of ≤ 0.05 followed by a post-hoc test (Scheffe) for significance. Significant transcripts were evaluated for relevance to canonical pathways and biological functions using Ingenuity Pathways Analysis (Ingenuity Systems, <http://www.ingenuity.com>). Heat maps were generated using Eisen Lab Cluster and Treeview software <http://rana.lbl.gov/EisenSoftware.htm>.

Gene Set Enrichment Analysis (GSEA; <http://www.broad.mit.edu/gsea/>) was used to evaluate whether a pre-defined set of genes showed statistically significant, concordant differences between two biological states

[23]. GSEA generates a list of genes from submitted .cel files and tests whether, within that queried list of genes, there is a statistically significant enrichment or not of pre-defined groups of genes, or “gene sets”. Gene sets examined through GSEA include canonical, metabolic and signaling pathways as well as groups of genes previously identified and validated to be up- or down-regulated when cells are given a particular stimulus. This type of analysis can sometimes detect more subtle changes present in the data. Expression profiles were submitted to GSEA using default settings and searched for enriched gene sets. Gene set C2 (curated gene sets) includes genes from online pathway databases, publications in PubMed, and knowledge from domain experts. Gene set C3 (motif gene sets) contains genes that share a transcription factor (TF) binding site defined in TRANSFAC[®], a database for transcription factors and their genomic binding sites. Additional .cel files were analyzed using GSEA including those from the livers of wild-type and PPAR α -null mice exposed to WY (400 μ L of a 10 mg/mL solution/day) for 6 hr or 5 days [24].

A description of the HS and WY exposure microarray experiment is available through Gene Expression Omnibus (GEO) at the National Center for Biotechnology Information <http://www.ncbi.nlm.nih.gov/geo/>, as accession number GSE14,869.

We categorized the genes regulated by HS in wild-type or PPAR α -null mice based on whether they were altered by HS in mouse embryonic fibroblasts (MEF) from wild-type or HSF1-null mice [8]. From this dataset, genes which fell into three groups were identified including 1) HSF1-dependent regulation by HS, 2) HSF1-independent regulation by HS and 3) altered regulation between wild-type and HSF1-null MEFs in the absence of HS. Genes were identified using procedures outlined in [25].

Real Time RT-PCR Analysis

The levels of gene expression were quantified using real time RT-PCR analysis. Real-time reverse transcriptase-PCR was performed as follows. After DNase treatment, total RNA was quantified (Ribogreen[®], Molecular Probes, Inc., Eugene, OR) and diluted with water. Fifty ng of RNA and PCR reagents were aliquoted into 96-well plates using an ABI Prism[™] 6700 Automated Nucleic Acid Workstation (Applied Biosystems, Foster City, CA) and subjected to real-time quantitative PCR (TaqMan[®], Applied Biosystems) using gene-specific primers (Additional File 1) and fluorescently labeled probes (Molecular Probes) designed by the Primer Express[®] software (Applied Biosystems). Amplification curves were generated using the ABI Prism[™] 7900 HT Sequence Detection System (Applied Biosystems). Expression relative to vehicle control animals was determined after normalizing to the ribosomal 18S gene. There were four animals

differences were caused by HS in wild-type mice and by WY+HS in PPAR α -null mice. Consistent with the PCA results, HS, WY alone or WY+HS treatments in wild-type mice altered 107, 1714 or 1418 genes, respectively (Figure 1B). Exposure to HS in wild-type mice altered a number of well-characterized targets of HSF1 (described below). Almost half of the HS regulated genes were also regulated by WY in the same direction. (The complete list of genes is found in Additional File 2.)

The gene expression pattern after treatment was dramatically different in PPAR α -null mice. HS, WY alone or WY+HS treatments in PPAR α -null mice altered 774, 6, or 68 genes, respectively (Figure 1B).

We directly compared the genes that were regulated by HS in the two strains (Figure 2A). A small number

of genes exhibited concordant expression changes in both strains including *Hsp90aa1*, *Hspe1*, and *Hspd1*. Other genes exhibited up-regulation in wild-type mice but down-regulation in PPAR α -null mice. There were 76 genes that exhibited altered expression only in wild-type mice, whereas 743 genes were altered only in PPAR α -null mice. For the most part, the HS genes unique to each strain lacked inclusion of Hsp family members.

The HS genes were categorized based on their expression behavior in the two strains after HS and WY exposure (Figure 2B). Genes in class I and II were altered by either HS alone (class I) or by HS and WY (class II) in a PPAR α -dependent manner. The class I genes included FK506 binding protein 4 (*Fkbp4*) and GrpE-like 2,




Figure 2 Heat shock alters the expression of different sets of genes in wild-type and PPAR α -null mice. **A**. Direct comparison of HS genes in wild-type and PPAR α -null mice. The relatively small number of genes which exhibited altered expression in both strains are shown in detail. **B**. Different classes of HS genes. The genes altered by HS were divided into six classes (I - VI) based on expression in wild-type and PPAR α -null mice and comparison with WY. The genes whose expression was confirmed by TaqMan are underlined. Red, up-regulation; green, down-regulation; black, no change. The intensity scales indicates fold-change due to exposure relative to controls. **C**. Altered expression of genes assessed by TaqMan. There were four animals per treatment group, and each sample was analyzed in duplicate. Variability is expressed as standard error of the mean. Means and S.E. ($n = 4$) for RT-PCR data were calculated by Student's t test. The level of significance was set at $p \leq 0.05$.

mitochondrial gene (*Grpel2*). Also included was ubiquitin carboxy-terminal hydrolase L1 (*Uchl1*) identified using less stringent microarray analysis cut-offs but confirmed by RT-PCR (Figure 2C). The class II genes included *Dnaja1* (*Hsp40*) and a tubulin family member (*Tubb2a*). RT-PCR revealed concordance with array expression for *Tubb2a*; *Dnaja1* exhibited weak induction by HS in PPAR α -null mice. Genes in class III were altered only by HS and were partially, if not completely independent of PPAR α , including stress-inducible phosphoprotein-1 (*Stip1*) in which increases in expression after HS in PPAR α -null mice were detected by RT-PCR. Genes in class IV were induced by HS independent of PPAR α and also exhibited PPAR α -dependent WY induction. These genes included a number of chaperones *Hsp90aa1*, *Hspe1*, *Hspa4l*, *Hspd1* and *Hsph1*. The expression pattern of *Hsp90aa1* and *Hsph1* was confirmed by RT-PCR. Class V genes were those that exhibited altered expression only in PPAR α -null mice after HS but not WY and included two metallothionein genes (*Mt1*, *Mt2*). Class VI genes were those that exhibited altered regulation after HS in PPAR α -null mice but not after HS+WY indicating suppression of HS transcriptional effects by WY. These genes included the up-regulation of serum amyloid protein family members (*Saa2*, *Saa3*) and down-regulation of a large number of mitochondrial genes (described below). Overall, the data is consistent with a subset of 1) HS-regulated genes (Class I and II) that are dependent on PPAR α for altered expression, 2) HS-regulated genes (Class III and IV) that are independent of PPAR α and 3) genes responsive to HS only in the absence of PPAR α (class V and VI).

Heat shock activates different biological responses in wild-type and PPAR α -null mice

We identified sets of genes that exhibited altered regulation in the two strains using Gene Set Enrichment Analysis (GSEA). GSEA is an analysis method that evaluates the expression of biological pathways on defined gene sets, as an alternative to examining individual genes, to assist in identifying significant biological changes in microarray data sets [23]. Consistent with HS effects on proteome maintenance genes, HS in wild-type mice increased the expression of genes involved in protein synthesis (HSA03010_RIBOSOME, RIBOSOMAL_PROTEINS, TRNA_SYNTHETASES) and degradation (HSA03050_PROTEASOME, PROTEASOME_DEGRADATION, PROTEASOMEPATHWAY, PROTEASOME) (Additional File 3). There were no gene sets that exhibited significant overlap with the down-regulated HS wild-type genes.

HS in PPAR α -null mice altered the expression of genes in a different set of biological categories (Additional File 4) including the up-regulation of genes associated with mouse liver tumors (LEE_DENA_UP,

LEE_MYC_E2F1_UP, LEE_E2F1_UP, LEE_MYC_TGFA_UP, LEE_ACOX1_UP, LEE_CIP_UP, LEE_MYC_TGFA_DN), muscle contraction (STRIATED_MUSCLE_CONTRACTION) and the aging kidney (AGEING_KIDNEY_UP).

There were a large number of genes down-regulated by HS in PPAR α -null mice associated with mitochondrial biogenesis and function regulated by the PPAR γ Co-activator 1 (PGC-1) family which includes PGC-1 α , PGC-1 β and PRC (MOOTHA_VOXPPOS, PGC, HUMAN_MITODDB_6_2002, MITOCHONDRIA) (Additional File 5). There was also down-regulation of genes involved in mouse and human liver tumors (LEE_MYC_E2F1_DN, LEE_DENA_DN, LEE_MYC_TGFA_DN, LEE_CIP_DN, HCC_SURVIVAL_GOOD_VS_POOR_UP) and down-regulation of genes associated with amino acid and fatty acid metabolism (VALINE_LEUCINE_AND_ISOLEUCINE_DEGRADATION, HSA00280_VALINE_LEUCINE_AND_ISOLEUCINE_DEGRADATION, HSA00380_TRYPTOPHAN_METABOLISM, HSA00071_FATTY_ACID_METABOLISM). The fatty acid metabolism genes included those that are regulated by PPAR α upon exposure to PPC (discussed below).

GSEA was also used to identify candidate transcription factors responsible for altered regulation of the HS genes. A number of stress-inducible transcription factors are likely involved in the altered expression of HS genes in wild-type mice. Consistent with well-characterized mechanisms of HS gene regulation, GSEA twice identified HSF1 as a candidate transcription factor of the HS genes (Additional File 6). Two sets of genes positively regulated by HS overlapped with those regulated by X-box binding protein (XBP1). XBP1 is a key component of the endoplasmic reticulum (ER) stress response [29]. Accumulation of unfolded proteins in the ER activates the unfolded protein response (UPR), resulting in transcriptional induction of ER chaperones and proteases in part through activation of XBP1. A number of genes associated with XBP1 were either up- (HOXC4, STAT3, PAK1IP1, SGK, VDP, SEC61A1, SFRS2, UBQLN1, MORF4L2) or down-regulated (COL1A2, ATBF1, POLG, GRIA3) in wild-type mice. HS genes linked to up-regulation of nuclear respiratory factor 1 (NRF1) were also identified (data not shown). NRF1 is activated by increased levels of heme and regulates a large number of genes involved in mitochondrial biogenesis and oxidative stress that overlap with those regulated by PGC-1 α or PGC-1 β [30,31].

In contrast, the genes regulated by HS in PPAR α -null mice overlapped with serum response factor (SRF) (one set significantly and three approaching significance) (Additional File 7). SRF is a ubiquitously expressed transcription factor that plays a role in regulating early response genes to various stimuli including healing after

tissue injury [32]. SRF plays a prominent role in regulating the expression of heart muscle genes [33] and may explain increases in muscle contraction genes (ACTA1, ACTA2, DES, MYL1, MYL2, MYL3, MYL9, MYOM1, TNNC2, TNNT1, TNNT3, TPM2, TPM4) after HS. Two gene sets that approached significance were regulated by HSF1. In summary, these results indicate that the PPAR α genotype has a dramatic effect on the biological pathways that are affected by HS.

PPAR α -independent changes in triglyceride levels and fatty acid oxidation protein expression by heat shock

We examined the levels of a number of clinical chemistry markers including triglycerides in the blood from mice 4 or 24 hrs after HS. The levels of alkaline phosphatase were increased in all WY-treated wild-type mice but not PPAR α -null mice (Figure 3A), paralleling the increases in gene expression for alkaline phosphatase, liver/bone/kidney (*Alpl*) after exposure to WY and other PPC (data not shown). Consistent with PPAR α agonists playing a role in glucose homeostasis [34,35], glucose levels were lower in wild-type but not PPAR α -null WY-treated mice at 4 and 24 hrs. HS also lowered glucose levels in wild-type but not PPAR α -null mice at 24 hrs. HS did not induce overt liver damage as measured by ALT levels; minor increases were observed by HS+WY treatment in wild-type but not PPAR α -null mice at 4 and 24 hrs that approached significance (data not

shown). These results were consistent with a lack of pathological findings in the liver. In wild-type mice triglyceride levels were increased by HS and decreased by WY at 4 hrs with no change by both treatments. At 24 hrs, triglyceride levels were decreased by HS, WY or HS+WY in wild-type mice and by HS or HS+WY in PPAR α -null mice. Despite uniform increases in cholesterol biosynthetic genes after WY exposure that were reversed by WY+HS treatment (Additional File 8 and 9), no changes in circulating cholesterol were noted between control and treated groups for each strain (data not shown).

Triglyceride levels are controlled in part by enzymes involved in fatty acid oxidation. Expression of fatty acid metabolism genes/proteins and triglyceride levels are reciprocally related. HS decreased the transcript levels of a number of genes involved in fatty acid transport and metabolism in both mouse strains including the *Acox1* gene (Figure 3B). In wild-type mice we observed the expected increases in enzymes involved in fatty acid oxidation including *Aco* and *Cyp4a* after exposure to WY in wild-type but not PPAR α -null mice 24 hrs after HS in liver (Figure 4A, B) and kidney (Figure 4C). HS had complex effects on expression of *Aco* and *Cyp4a*. In the liver HS decreased the expression of ACO-B protein in wild-type mice at 24 hrs. In the kidney but not the liver, ACO-B protein was increased by HS in PPAR α -





Figure 3 PPAR α -independent changes in triglyceride levels and fatty acid metabolism genes by heat shock. A. Serum levels of alkaline phosphatase, glucose and triglycerides 4 and 24 hrs after HS. B. Decreases in fatty acid metabolism genes by HS in wild-type and PPAR α -null mice. Genes called as significantly altered between control and HS groups were extracted from GSEA gene sets HSA03320_PPAR_SIGNALING_PATHWAY (wild-type mice) and HSA00071_FATTY_ACID_METABOLISM (PPAR α -null mice).

null mice. In the liver but not the kidney, *Cyp4a* protein was induced by HS only in PPAR α -null mice. HS-inducible gene expression of *Cyp4a10/14* genes was not observed by microarray indicating post-transcriptional regulation of expression.

These findings indicate that HS elicits complex changes in fatty acid metabolism gene and protein expression leading to alterations in triglyceride levels, some of which are independent of PPAR α . The increases in triglyceride levels in wild-type mice by HS at 4 hrs may be attributed in part to decreases in fatty acid oxidation genes; the decreases in triglyceride levels in PPAR α -null mice are correlated with increases in *Cyp4a* and ACO-B in the liver or kidney. The decreases in triglyceride levels in wild-type mice 24 hrs after HS cannot be explained by the decreases in ACO and *Cyp4a* in liver and kidney. Thus, additional mechanisms that control triglyceride levels are likely at play including effects on mitochondrial fatty acid oxidation and require further study. In summary, 24 hrs after HS the decreases in triglyceride levels were PPAR α -independent whereas decreases in glucose levels were PPAR α -dependent.

Chaperone protein expression after heat shock and WY treatment

We examined protein expression of Hsp family members in livers and kidneys of the mice 24 hrs after HS. Given the transcriptional increases in chaperonin-containing T-complex 1 (Tcp-1) family members *Cct3*, *Cct4*, *Cct7* and *Cct8* after exposure to WY in wild-type mice (Additional File 2), we also examined the expression of Tcp1 η protein. Western blots consistently detected two bands of ~60 kDa (full-length Tcp1 η) and an immunoreactive fragment of ~40 kDa (Tcp1 η (40)). Increases in Tcp1 η (40), Hsp25, Hsp70, Hsp86, Hsp110 but not Hsp65 were observed in wild-type mice after HS; the increases in all of the proteins were PPAR α -independent, as increases were also observed in HS-treated PPAR α -null mice (Figure 5A, B, C). In contrast, TCP1 η exhibited decreased expression after HS in wild-type and PPAR α -null mice. WY exposure led to increased expression in Tcp1 η , Hsp70, Hsp86, and Hsp110 in wild-type mice that was partially or completely eliminated in PPAR α -null mice. The increases in protein expression after HS and WY were generally






Figure 5 Expression of chaperone proteins after heat shock or PPC treatment. A. Expression of Hsp in the livers of mice. Proteins were extracted from the livers of wild-type or PPAR α -null mice on control or WY diet, 24 hrs after HS or mock HS. Expression was assessed by Western blot using primary antibodies against the indicated proteins. B and C. Quantitation of the western blots in A. D. Altered protein expression in the kidneys of wild-type and PPAR α -null mice exposed to HS or WY. Protein expression was assessed in the kidneys of the mice described in A.

consistent with the microarray and RT-PCR data for *Hspa1a* and *Hspa1b* (Hsp70), *Hsp90aa1* (Hsp86/90), *Hsph1* (Hsp105/110) and *Cct7* (TCP1 η). No transcriptional changes by microarray were detected for *Hspb1* (Hsp25), but RT-PCR showed increases after a 7 day WY exposure in a previous study [12].

We assessed protein expression in the kidneys from the animals described in this study. Tcpl η , although not affected by HS in liver was induced up to 8-fold by HS in kidneys from both strains, indicating tissue-specific induction by HS (Figure 5D). Hsp86 but not Hsp25, Hsp70, and Hsp110 was induced by HS in the kidney (Figure 5D and data not shown). After WY exposure Hsp86, but not Tcpl η , Hsp25, Hsp70, or Hsp110 was weakly induced in the kidneys of wild-type, but not PPAR α -null mice.

Most of the Hsp genes/proteins that were induced by WY (Hsp25, Hsp70, Hsp86, Hsp110, TCP1) carry out protein folding in the cytoplasm in a coordinated manner. For example, Hsp70 and Hsp110 cooperate to

sequentially fold proteins in the cytoplasm [36]. In contrast, we did not observe any mitochondrial chaperones such as mtHsp70 altered on the transcriptional level; a number of chaperones which operate in the endoplasmic reticulum including BiP/Grp78 [37] and Grp94 (our study) were down-regulated. An earlier report showed that the Grp94 gene and protein were increased after exposure to WY or nafenopin [38]. Overall, our studies indicate that WY coordinately regulates a number of chaperones which operate in the cytoplasm.

What might the increased levels of chaperone proteins be doing in the liver after WY exposure? Increased levels of chaperones might allow tight control of the inducibility of PPAR α . Many nuclear receptors interact with chaperone proteins including the ones induced by WY in our studies [39]. PPAR α interacts with Hsp72 [40] and is inhibited by Hsp90 [41]. Thus, induction of Hsp90 and other family members may dampen the PPAR α transcriptional response. Additionally, Hsp induction may help support the increases in protein

synthesis associated with hepatocyte replication after PPC exposure. In addition, increased expression of Hsp family members has been associated with protection from apoptosis [42,43] and PPC, at least under acute exposure conditions decrease basal levels of apoptosis [10]. PPC exposure leads to dramatic increases in peroxisome size and number and increased expression of TCP1 subunits may be important for proper protein insertion into the peroxisomal membrane [44].

It is generally appreciated that increased expression of Hsps by mild HS can protect from subsequent challenges by more severe treatments [45]. Our microarray and protein expression analysis showed that WY can induce the gene and protein expression of a number of chaperones. These same chaperones including Hsp25/27 [46,47], Hsp70 [45,48], Hsp86/90 [49-51] and Hsp105/110 [36,52,53] when over-expressed can protect cells from various stressors. Our results identify a number of potential candidate genes/proteins that could protect cells from environmental stressors by pretreatment with PPAR α agonists.

Global decreases in mitochondrial gene expression in PPAR α -null mice after heat shock are prevented by WY pretreatment

HS can induce mitochondrial swelling, loss of mitochondria and uncoupling of oxidative phosphorylation [45]. HS in PPAR α -null mice led to dramatic decreases in mitochondrial biogenesis and mitochondrial oxidative phosphorylation genes regulated by PGC-1 family members (Additional File 6). Coordinated down-regulation of a curated set of genes regulated by PGC-1 was observed; out of the 310 genes in this list shared by the U74Av2 array, 123 were down-regulated (Figure 6A). Pretreatment of PPAR α -null mice with WY led to resistance to gene expression changes, as far fewer genes were altered by both treatments compared to HS alone (774 genes for HS alone vs. 68 genes for HS+WY). We compared the two groups directly using GSEA and identified significant increases in electron transporter genes after HS +WY treatment compared to HS alone including those that function in the mitochondria (Figure 6B). Significant alterations in mitochondrial gene sets were not observed in a comparison between control and HS+WY groups using GSEA (data not shown). Thus, WY pretreatment of PPAR α -null mice prevented decreases in mitochondrial gene expression due to HS.

To identify candidates involved in resistance to HS-induced decreases in mitochondrial gene expression, we searched for genes regulated by WY independently of PPAR α . Our previous experiments indicated that WY had minimal effects in the livers of PPAR α -null mice, altering only up to ~8% of the total number of genes compared to wild-type mice [54]. These results were based on transcript profiles generated using the

Affymetrix gene chip U74Av2 which contains ~12,000 genes. Using full-genome arrays containing ~45,000 genes (Affymetrix MOE430_2 arrays), we examined gene expression in PPAR α -null mice after exposure to WY for 6 hrs or 5 days. WY had minimal effects on gene expression after 6 hrs in PPAR α -null mice (1 gene change) but at 5 days, 92 genes were significantly altered in PPAR α -null mice. A number of the changes would affect intermediary metabolism and mitochondrial gene expression, the most striking of these changes being increases in PGC-1 β (*Ppargc1b*) (~2-fold, $p = 0.01$; Figure 6C). (This gene is not included on the U74Av2 array.) Genes known to be regulated by PGC-1 β included *Cpt1* [55-57] and a number of others involved in fatty acid oxidation including acetyl-Coenzyme A acyltransferase 1B (*Acaa1b*), aldehyde dehydrogenase family 3, subfamily A2 (*Aldh3a2*), acetyl-Coenzyme A acyltransferase 1A (*Acaa1a*), enoyl coenzyme A hydratase 1, peroxisomal (*Ech1*) and peroxisomal delta3, delta2-enoyl-Coenzyme A isomerase (*Peci*) (Figure 6D).

PGC-1 β is a master controller of hepatic energy homeostasis in part through regulation of mitochondrial oxidative metabolism genes. Overproduction of PGC-1 β leads to increases in mitochondrial volume density in C2C12 myotubes [58], and defects in PGC-1 β expression have been associated with global decreases in mitochondrial gene expression in the skeletal muscle of diabetics [30]. PGC-1 β physically interacts with transcription factors involved in mitochondrial biogenesis and lipid metabolism including Nrf-1, PPAR α and PPAR γ [55,59,60]. In trans-activation assays, WY activates PPAR β and PPAR γ , albeit to a lesser extent than PPAR α [61,62]. Increases in PGC-1 β co-activator function in the presence of the weak agonist function of WY may lead to increased expression of fatty acid oxidation genes regulated by PPAR β or PPAR γ similar to what we have observed with the PPC perfluorooctanoic acid in PPAR α -null mice [54]. These results indicate that WY can alter the expression and activity of effectors of mitochondrial biogenesis and fatty acid oxidation, preventing HS from decreasing those gene batteries in PPAR α -null mice. This protective effect of WY is reminiscent of thermotolerance induced by prior treatment with a low temperature HS [45].

Hsp gene regulation by PPAR α and HSF1

HSF1 controls expression of *Hsp* genes after HS [7,8]. We determined if there was any overlap in HS genes regulated by HSF1 and PPAR α . We identified a number of classes of HSF1-dependent or -independent genes regulated by HS as well as genes that require HSF1 for basal expression (Additional File 10 and 11). We then identified genes that overlapped with those regulated by HS in wild-type and PPAR α -null mice from our study. The expression of the 45 unique pairs of gene identifiers from each array is shown in Figure 7A.




Figure 6 Global decreases in mitochondrial gene expression in PPAR α -null mice after heat shock are prevented by WY pretreatment.

A. Down-regulation of genes regulated by the co-activator PGC-1. Gene Set Enrichment Analysis (GSEA) was used to identify gene sets that exhibited significant overlaps with those gene differences between control and HS in PPAR α -null mice. Left, enrichment plot for genes regulated by PGC-1. Black bars illustrate the position of probe sets belonging to the PGC-1 gene set in the context of all probes on the U74Av2 array. The running enrichment score (RES) plotted as a function of the position within the ranked list of array probes is shown as a green line. The ranked list metric shown in gray illustrates the correlation between the signal to noise values of all individually ranked genes according to the class labels (experimental conditions). Right, individual expression profiles for leading edge probe sets contributing to the normalized enrichment score are shown. Signal intensities are illustrated by varying shades of red (up-regulation) and blue (down-regulation). B. Prevention of down-regulation of electron transporter gene expression by pretreatment with WY. Left, enrichment plot for genes with electron transporter activity as described in A. Right, individual expression profiles for probe sets contributing to the normalized enrichment score are shown. C. Increased expression of PGC1 β and regulated genes involved in lipid homeostasis by WY in PPAR α -null mice. GSEA-derived heat map of the top 100 differentially expressed probe sets enriched in the control or WY-treated groups from PPAR α -null mice. Location of PGC1 β (*Ppargc1b*) and regulated genes are indicated. D. Increased expression of genes involved in fatty acid metabolism after exposure to WY for 5 days in PPAR α -null mice. Genes significantly altered in the GSEA set "fatty acid metabolism" are shown. These include a number that overlap with those that are involved in mitochondrial fatty acid metabolism regulated by PGC-1.

The genes fell into 4 groups based on dependence on PPAR α and HSF1 (Figure 7B). The largest groups were those that were PPAR α -independent and were about equally split between those that were dependent or independent of HSF1. PPAR α -dependent genes included those that were HSF1-dependent (*Cacybp*, *Dnaja1*, *Fkbp4*) or independent (*Apoe*, *Irak1*). This comparison reveals that although some genes were altered by HS independent of both PPAR α and HSF1, most genes required one or both factors for HS regulation.

An important question is why does PPAR α expression/activation determine in part the profile of HS genes? We hypothesize that PPAR α is required for the induction of a subset of the HS genes. Class I and II genes (Figure 2B) require PPAR α for HS regulation indicating that they could possess functional PPRES. Genes have been identified that possess HSEs that bind HSF1 but that were not HS-inducible. This provides evidence that other cis-acting elements or interacting factors were required for HS inducibility [8]. A linkage




Figure 7 Regulation of heat shock genes by HSF1 and PPARα. Genes which exhibited significant changes in expression due to HS from the dataset of Trinklein et al. (2004) and from the present study were identified. A. Heat map of gene expression changes by HS in wild-type (W) and PPARα-null (N) mice compared to the Trinklein et al. (2004) dataset. In the Trinklein et al. (2004) study, mouse embryonic fibroblasts were subjected to HS followed by recovery for the indicated times in hrs. Genes were subjected to one-dimensional hierarchical clustering. Red, up-regulation; green, down-regulation; grey, no data; black, no change. B. Classification of genes based on regulation by PPARα and HSF1.

between HSE and PPRES in the promoters of HS genes was found as binding sites for PPAR family members were significantly increased within the 1000 bp upstream of the genes that were up- or down-regulated by HS in a human epithelial cell line [63]. Follow-up PPAR-HSF1 co-immunoprecipitation studies would reveal the nature of the interactions. Our analysis demonstrated major differences in the target genes of HS between wild-type and PPARα-null strains indicating that in the absence of PPARα, major changes in chromatin accessibility to HSF1 and other factors leads to greater sensitivity to HS alteration.

How are the HS genes regulated by PPC through PPARα? Given that *Hsp* gene expression is controlled in part by heat shock factor 1 (HSF1), one possibility is that the increases in *Hsp* gene expression are secondary to increases in the expression and activity of HSF1. However, we did not observe changes in *Hsf1* expression in our transcript profiling studies and earlier studies showed that HSF1 and HSF2 expression and binding to HSE was not altered by WY exposure in the rat liver [64]. To help determine whether regulation of *Hsp* gene expression is direct or indirect, we examined their promoters and

found that only a few genes possess a putative PPRES (data not shown) indicating indirect activation.

Many *Hsp* genes may be regulated indirectly through increases in oxidative stress. There is abundant evidence for the increased expression of chaperone gene expression under conditions that also increase oxidative stress [5,45]. PPC exposure leads to increases in oxidative stress and lipid peroxidation mediated through increased activities of enzymes that generate reactive oxygen species [10]. Although direct evidence that *Hsp* genes are activated by PPC-mediated increases in oxidative stress is lacking, the absolute increases in expression of some *Hsp* genes was higher in mice nullizygous for Nrf2, a transcription factor activated by oxidative stress that regulates genes that decrease oxidative stress. Thus, in the absence of Nrf2, increased levels of oxidative stress may have contributed to the greater increases in the *Hsp* genes by WY [12]. However, we cannot rule out the induction of a HSP1 cofactor secondary to PPARα induction." In summary, PPARα may regulate the *Hsp* genes in a species-specific manner secondarily to increases in oxidative stress. Further work is needed to confirm this hypothesis.

Conclusions

Our microarray analysis of HS in the livers of mice uncovered a dramatic influence of PPAR α on the overall gene expression pattern. There were remarkable differences in the transcriptional response to HS between wild-type and PPAR α -null mice. A number of HS genes required PPAR α for induction. Many mitochondrial genes were uniquely down-regulated in PPAR α -null but not wild-type mice. Pretreatment of PPAR α -null mice with WY prevented the down-regulation of these genes possibly through the increased expression of PGC-1 β and regulated genes. Our findings support the hypothesis that PPAR α in the liver acts as an integrator of chemical and physical stress signals manifested in the regulation of stress gene responses in the liver.

Additional file 1: Sequences of primers. Sequences of primers used in TaqMan studies.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S1.DOC>]

Additional file 2: Gene expression changes after WY, heat shock or heat shock+WY in wild-type and PPAR α -null mouse livers. Table represents the gene annotations and fold changes for gene expression changes after WY, heat shock or heat shock+WY in wild-type and PPAR α -null mouse livers.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S2.XLS>]

Additional file 3: Table of genesets significantly up-regulated by heat shock in wild-type mice. Table describes GSEA genesets significantly up-regulated by heat shock in wild-type mice.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S3.DOC>]

Additional file 4: Table of genesets significantly up-regulated by heat shock in PPAR α -null mice. Table describes GSEA genesets significantly up-regulated by heat shock in PPAR α -null mice.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S4.DOC>]

Additional file 5: Table of genesets significantly down-regulated by heat shock in PPAR α -null mice. Table describes GSEA genesets significantly down-regulated by heat shock in PPAR α -null mice.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S5.DOC>]

Additional file 6: Table of transcription factor genesets significantly up-regulated by heat shock in wild-type mice. Table describes the GSEA transcription factor genesets significantly up-regulated by heat shock in wild-type mice.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S6.DOC>]

Additional file 7: Table of transcription factor genesets significantly up-regulated by heat shock in PPAR α -null mice. Table describes the GSEA transcription factor genesets significantly up-regulated by heat shock in PPAR α -null mice.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S7.DOC>]

Additional file 8: Analysis of interactions between WY and HS in wild-type mice. Interactions between WY and HS in wild-type mice using Ingenuity Pathway Analysis Tool.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S8.DOC>]

Additional file 9: Canonical pathways analysis using Ingenuity.

Canonical pathways in an Ingenuity comparison between genes only regulated by WY or only by WY+HS in wild-type mice.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S9.XLS>]

Additional file 10: Excel spreadsheet of genes regulated by heat shock or by strain differences between wild-type and HSF1-null mice.

Genes were originally from the Trinklein et al. (2004) study. Excel spreadsheet of genes regulated by heat shock or by strain differences between wild-type and HSF1-null mice.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S10.XLS>]

Additional file 11: Figures of expression of heat shock genes in wild-type and HSF1-null mouse embryonic fibroblasts.

Mouse embryonic fibroblasts were given a HS and cells were harvested at the indicated times as described (Trinklein et al., 2004). Genes which exhibited significant changes after HS or that exhibited significant differences between wild-type and HSF1-null strains were identified as described in the Methods. A. HSF1-dependent HS genes. B. HSF1-independent HS genes. C. Genes which exhibited differences in expression between control wild-type and control HSF1-null strains.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-11-16-S11.PPT>]

Abbreviations

DEHP: di-(2-ethylhexyl)phthalate; HSE: heat shock elements; HSF: heat shock factor; HSP: heat shock proteins; PCR: polymerase chain reaction; PPAR: peroxisome proliferator-activated receptor; PPC: peroxisome proliferator chemical; PPRE: peroxisome proliferator response element; WY: WY-14,643.

Acknowledgements

We thank Drs. David Dix and Kevin Kregel for guidance in performing the animal exposures and western blots, Mr. Artie James, the CIIT Animal Care and Necropsy and Histology Units for assistance in performing these studies, Dr. Dennis House for assistance with statistical analyses, Mr. Lawrence Yoon for performing the RT-PCR analysis, Mr. John Havel for graphics support and Drs. David Dix, Stephen Edwards and Susan Hester for critical review of the manuscript.

The information in this document has been funded in part by the U.S. Environmental Protection Agency. It has been subjected to review by the National Health and Environmental Effects Research Laboratory and approved for publication. Approval does not signify that the contents reflect the views of the Agency, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

Author details

¹NHEERL Toxicogenomics Core, US EPA, Research Triangle Park, NC 27711, USA. ²Safety Assessment, Meril, Ltd., Duluth, GA 30096, USA. ³Department of Pharmacology, Physiology and Therapeutics, University of North Dakota, School of Medicine, 501 N. Columbia Road, Grand Forks, ND 58203-2817, USA. ⁴Nutrition, Metabolism and Genomics Group, Wageningen University and Nutrigenomics Consortium, TI Food and Nutrition, Wageningen, the Netherlands. ⁵Institute of Chemical and Engineering Sciences, A*STAR (Agency for Science, Technology and Research), 1 Pesek Road, Jurong Island, Singapore. ⁶Department of Chemical and Biomolecular Engineering, National University of Singapore, 10 Kent Ridge Crescent, Singapore.

Authors' contributions

BV analyzed the microarray and clinical chemistry data and helped to draft the manuscript. SA generated and analyzed the microarray data and analyzed the RT-PCR data. HBB analyzed the western data. HR analyzed the microarray data. SK generated the microarray data. SJ and RS analyzed the MEF microarray dataset. JCC conceived of the study, participated in study design and animal studies, generated western data, analyzed microarray data and helped to draft the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 17 February 2009

Accepted: 7 January 2010 Published: 7 January 2010

References

- Christians ES, Benjamin IJ: **Heat shock response: lessons from mouse knockouts.** *Handbook of experimental pharmacology* 2006, **172**: 139-152.
- Pirkkala L, Nykanen P, Sistonen L: **Roles of the heat shock transcription factors in regulation of the heat shock response and beyond.** *Faseb J* 2001, **15**(7):1118-1131.
- Shamovsky I, Nudler E: **New insights into the mechanism of heat shock response activation.** *Cell Mol Life Sci* 2008, **65**(6):855-861.
- Slavotinek AM, Biesecker LG: **Unfolding the role of chaperones and chaperonins in human disease.** *Trends Genet* 2001, **17**(9):528-535.
- Akerfelt M, Trouillet D, Mezger V, Sistonen L: **Heat shock factors at a crossroad between stress and development.** *Annals of the New York Academy of Sciences* 2007, **1113**:15-27.
- Arya R, Mallik M, Lakhotia SC: **Heat shock genes - integrating cell survival and death.** *Journal of biosciences* 2007, **32**(3):595-610.
- Page TJ, Sikder D, Yang L, Pluta L, Wolfinger RD, Kodadek T, Thomas RS: **Genome-wide analysis of human HSF1 signaling reveals a transcriptional program linked to cellular adaptation and survival.** *Molecular bioSystems* 2006, **2**(12):627-639.
- Trinklein ND, Murray JI, Hartman SJ, Botstein D, Myers RM: **The role of heat shock transcription factor 1 in the genome-wide regulation of the mammalian heat shock response.** *Molecular biology of the cell* 2004, **15**(3):1254-1261.
- Corton JC, Lapinskas PJ, Gonzalez FJ: **Central role of PPARalpha in the mechanism of action of hepatocarcinogenic peroxisome proliferators.** *Mutation research* 2000, **448**(2):139-151.
- Klaunig JE, Babich MA, Baetcke KP, Cook JC, Corton JC, David RM, DeLuca JG, Lai DY, McKee RH, Peters JM, et al: **PPARalpha agonist-induced rodent tumors: modes of action and human relevance.** *Critical reviews in toxicology* 2003, **33**(6):655-780.
- Anderson SP, Dunn C, Laughter A, Yoon L, Swanson C, Stulnig TM, Steffensen KR, Chandraratna RA, Gustafsson JA, Corton JC: **Overlapping transcriptional programs regulated by the nuclear receptors peroxisome proliferator-activated receptor alpha, retinoid x receptor, and liver x receptor in mouse liver.** *Molecular pharmacology* 2004, **66**(6):1440-1452.
- Anderson SP, Howroyd P, Liu J, Qian X, Bahnemann R, Swanson C, Kwak MK, Kensler TW, Corton JC: **The transcriptional response to a peroxisome proliferator-activated receptor alpha agonist includes increased expression of proteome maintenance genes.** *The Journal of biological chemistry* 2004, **279**(50):52390-52398.
- Peraza MA, Burdick AD, Marin HE, Gonzalez FJ, Peters JM: **The toxicology of ligands for peroxisome proliferator-activated receptors (PPAR).** *Toxicol Sci* 2006, **90**(2):269-295.
- Corton JC, Brown-Borg HM: **Peroxisome proliferator-activated receptor gamma coactivator 1 in caloric restriction and other models of longevity.** *The journals of gerontology* 2005, **60**(12):1494-1509.
- Mehendale HM: **PPAR-alpha: a key to the mechanism of hepatoprotection by clofibrate.** *Toxicol Sci* 2000, **57**(2):187-190.
- Chen C, Hennig GE, Whiteley HE, Corton JC, Manautou JE: **Peroxisome proliferator-activated receptor alpha-null mice lack resistance to acetaminophen hepatotoxicity following clofibrate exposure.** *Toxicol Sci* 2000, **57**(2):338-344.
- Corton JC, Apte U, Anderson SP, Limaye P, Yoon L, Latendresse J, Dunn C, Everitt JJ, Voss KA, Swanson C, et al: **Mimetics of caloric restriction include agonists of lipid-activated nuclear receptors.** *The Journal of biological chemistry* 2004, **279**(44):46204-46212.
- Anderson SP, Yoon L, Richard EB, Dunn CS, Cattley RC, Corton JC: **Delayed liver regeneration in peroxisome proliferator-activated receptor-alpha-null mice.** *Hepatology (Baltimore, Md)* 2002, **36**(3):544-554.
- Portilla D, Dai G, Peters JM, Gonzalez FJ, Crew MD, Proia AD: **Etomoxir-induced PPARalpha-modulated enzymes protect during acute renal failure.** *American journal of physiology* 2000, **278**(4):F667-675.
- Sivarajah A, Chatterjee PK, Hattori Y, Brown PA, Stewart KN, Todorovic Z, Mota-Filipe H, Thiemermann C: **Agonists of peroxisome-proliferator activated receptor-alpha (clofibrate and WY14643) reduce renal ischemia/reperfusion injury in the rat.** *Med Sci Monit* 2002, **8**(12):BR532-539.
- Lee SS, Pineau T, Drago J, Lee EJ, Owens JW, Kroetz DL, Fernandez-Salguero PM, Westphal H, Gonzalez FJ: **Targeted disruption of the alpha isoform of the peroxisome proliferator-activated receptor gene in mice results in abolishment of the pleiotropic effects of peroxisome proliferators.** *Molecular and cellular biology* 1995, **15**(6):3012-3022.
- Wilson CL, Miller CJ: **Simpleaffy: a BioConductor package for Affymetrix Quality Control and data analysis.** *Bioinformatics (Oxford, England)* 2005, **21**(18):3683-3685.
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, et al: **Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.** *Proceedings of the National Academy of Sciences of the United States of America* 2005, **102**(43):15545-15550.
- Sanderson LM, de Groot PJ, Hooiveld GJ, Koppen A, Kalkhoven E, Muller M, Kersten S: **Effect of synthetic dietary triglycerides: a novel research paradigm for nutrigenomics.** *PLoS ONE* 2008, **3**(2):e1681.
- Jonnalagadda S, Srinivasan R: **Principal components analysis based methodology to identify differentially expressed genes in time-course microarray data.** *BMC bioinformatics* 2008, **9**:267.
- Fan LQ, Coley J, Miller RT, Cattley RC, Corton JC: **Opposing mechanisms of NADPH-cytochrome P450 oxidoreductase regulation by peroxisome proliferators.** *Biochemical pharmacology* 2003, **65**(6):949-959.
- Corton JC, Fan LQ, Brown S, Anderson SP, Bocos C, Cattley RC, Mode A, Gustafsson JA: **Down-regulation of cytochrome P450 2C family members and positive acute-phase response gene expression by peroxisome proliferator chemicals.** *Molecular pharmacology* 1998, **54**(3):463-473.
- Dinh HK, Zhao B, Schuschereba ST, Merrill G, Bowman PD: **Gene expression profiling of the response to thermal injury in human cells.** *Physiological genomics* 2001, **7**(1):3-13.
- Yoshida H: **Unconventional splicing of XBP-1 mRNA in the unfolded protein response.** *Antioxidants & redox signaling* 2007, **9**(12):2323-2333.
- Patti ME, Butte AJ, Crunkhorn S, Cusi K, Berria R, Kashyap S, Miyazaki Y, Kohane I, Costello M, Saccone R, et al: **Coordinated reduction of genes of oxidative metabolism in humans with insulin resistance and diabetes: Potential role of PGC1 and NRF1.** *Proceedings of the National Academy of Sciences of the United States of America* 2003, **100**(14):8466-8471.
- Spiegelman BM: **Transcriptional control of energy homeostasis through the PGC1 coactivators.** *Novartis Foundation symposium* 2007, **286**:3-6, discussion 6-12, 162-163, 196-203.
- Chai J, Tarnawski AS: **Serum response factor: discovery, biochemistry, biological roles and implications for tissue injury healing.** *J Physiol Pharmacol* 2002, **53**(2):147-157.
- Miano JM, Long X, Fujiwara K: **Serum response factor: master regulator of the actin cytoskeleton and contractile apparatus.** *Am J Physiol Cell Physiol* 2007, **292**(1):C70-81.
- Cho MC, Lee K, Paik SG, Yoon DY: **Peroxisome Proliferators-Activated Receptor (PPAR) Modulators and Metabolic Disorders.** *PPAR research* 2008, **2008**:679137.
- Fievat C, Fruchart JC, Staels B: **PPARalpha and PPARgamma dual agonists for the treatment of type 2 diabetes and the metabolic syndrome.** *Current opinion in pharmacology* 2006, **6**(6):606-614.
- Easton DP, Kaneko Y, Subjeck JR: **The hsp110 and Grp1 70 stress proteins: newly recognized relatives of the Hsp70s.** *Cell stress & chaperones* 2000, **5**(4):276-290.
- Motojima K, Goto S: **Rat liver BIP/GRP78 is down-regulated by a peroxisome-proliferator, clofibrate.** *FEBS letters* 1992, **308**(2):207-210.
- Macdonald N, Barrow K, Tonge R, Davison M, Roberts RA, Chevalier S: **PPARalpha-dependent alteration of GRP94 expression in mouse**

- hepatocytes. *Biochemical and biophysical research communications* 2000, **277**(3):699-704.
39. Morimoto Ri: **Dynamic remodeling of transcription complexes by molecular chaperones.** *Cell* 2002, **110**(3):281-284.
 40. Huang Q, Alvares K, Chu R, Bradfield CA, Reddy JK: **Association of peroxisome proliferator-activated receptor and Hsp72.** *The Journal of biological chemistry* 1994, **269**(11):8493-8497.
 41. Sumanasekera WK, Tien ES, Davis JW, Turpey R, Perdew GH, Vanden Heuvel JP: **Heat shock protein-90 (Hsp90) acts as a repressor of peroxisome proliferator-activated receptor-alpha (PPARalpha) and PPARbeta activity.** *Biochemistry* 2003, **42**(36):10726-10735.
 42. Beere HM: **Stressed to death: regulation of apoptotic signaling pathways by the heat shock proteins.** *Sci STKE* 2001, **2001**(93):RE1.
 43. Latchman DS: **Heat shock proteins and cardiac protection.** *Cardiovascular research* 2001, **51**(4):637-646.
 44. Pause B, Diestelkotter P, Heid H, Just WW: **Cytosolic factors mediate protein insertion into the peroxisomal membrane.** *FEBS letters* 1997, **414**(1):95-98.
 45. Benjamin UJ, McMillan DR: **Stress (heat shock) proteins: molecular chaperones in cardiovascular biology and disease.** *Circulation research* 1998, **83**(2):117-132.
 46. Martin JL, Mestrlil R, Hilal-Dandan R, Brunton LL, Dillmann WH: **Small heat shock proteins and protection against ischemic injury in cardiac myocytes.** *Circulation* 1997, **96**(12):4343-4348.
 47. Park YM, Han MY, Blackburn RV, Lee YJ: **Overexpression of HSP25 reduces the level of TNF alpha-induced oxidative DNA damage biomarker, 8-hydroxy-2'-deoxyguanosine, in L929 cells.** *Journal of cellular physiology* 1998, **174**(1):27-34.
 48. Tolson JK, Dix DJ, Voellmy RW, Roberts SM: **Increased hepatotoxicity of acetaminophen in Hsp70i knockout mice.** *Toxicology and applied pharmacology* 2006, **210**(1-2):157-162.
 49. Conconi M, Petropoulos I, Emod I, Turlin E, Biville F, Friguet B: **Protection from oxidative inactivation of the 20S proteasome by heat-shock protein 90.** *The Biochemical journal* 1998, **333**(Pt 2):407-415.
 50. Cumming DV, Heads RJ, Watson A, Latchman DS, Yellon DM: **Differential protection of primary rat cardiocytes by transfection of specific heat stress proteins.** *Journal of molecular and cellular cardiology* 1996, **28**(12):2343-2349.
 51. Lee MW, Park SC, Chae HS, Bach JH, Lee HJ, Lee SH, Kang YK, Kim KY, Lee WB, Kim SS: **The protective role of HSP90 against 3-hydroxykynurenine-induced neuronal apoptosis.** *Biochemical and biophysical research communications* 2001, **284**(2):261-267.
 52. Hatayama T, Yamagishi N, Minobe E, Sakai K: **Role of hsp105 in protection against stress-induced apoptosis in neuronal PC12 cells.** *Biochemical and biophysical research communications* 2001, **288**(3):528-534.
 53. Oh HJ, Chen X, Subjeck JR: **Hsp110 protects heat-denatured proteins and confers cellular thermoresistance.** *The Journal of biological chemistry* 1997, **272**(50):31636-31640.
 54. Rosen MB, Lee JS, Ren H, Vallanat B, Liu J, Waalkes MP, Abbott BD, Lau C, Corton JC: **Toxicogenomic dissection of the perfluorooctanoic acid transcript profile in mouse liver: evidence for the involvement of nuclear receptors PPAR alpha and CAR.** *Toxicol Sci* 2008, **103**(1):46-56.
 55. Lin J, Tarr PT, Yang R, Rhee J, Puigserver P, Newgard CB, Spiegelman BM: **PGC-1beta in the regulation of hepatic glucose and energy metabolism.** *The Journal of biological chemistry* 2003, **278**(33):30843-30848.
 56. Sadana P, Zhang Y, Song S, Cook GA, Elam MB, Park EA: **Regulation of carnitine palmitoyltransferase I (CPT-1alpha) gene expression by the peroxisome proliferator activated receptor gamma coactivator (PGC-1) isoforms.** *Molecular and cellular endocrinology* 2007, **267**(1-2):6-16.
 57. Wolfrum C, Stoffel M: **Coactivation of Foxa2 through Pgc-1beta promotes liver fatty acid oxidation and triglyceride/VLDL secretion.** *Cell metabolism* 2006, **3**(2):99-110.
 58. St-Pierre J, Lin J, Krauss S, Tarr PT, Yang R, Newgard CB, Spiegelman BM: **Bioenergetic analysis of peroxisome proliferator-activated receptor gamma coactivators 1alpha and 1beta (PGC-1alpha and PGC-1beta) in muscle cells.** *The Journal of biological chemistry* 2003, **278**(29):26597-26603.
 59. Kressler D, Schreiber SN, Knutti D, Kralli A: **The PGC-1-related protein PERC is a selective coactivator of estrogen receptor alpha.** *The Journal of biological chemistry* 2002, **277**(16):13918-13925.
 60. Lin J, Puigserver P, Donovan J, Tarr P, Spiegelman BM: **Peroxisome proliferator-activated receptor gamma coactivator 1beta (PGC-1beta), a novel PGC-1-related transcription coactivator associated with host cell factor.** *The Journal of biological chemistry* 2002, **277**(3):1645-1648.
 61. Corton JC, Lapinskas PJ: **Peroxisome proliferator-activated receptors: mediators of phthalate ester-induced effects in the male reproductive tract?** *Toxicol Sci* 2005, **83**(1):4-17.
 62. Lapinskas PJ, Brown S, Leesnitzer LM, Blanchard S, Swanson C, Cattley RC, Corton JC: **Role of PPARalpha in mediating the effects of phthalates and metabolites in the liver.** *Toxicology* 2005, **207**(1):149-163.
 63. Laramie JM, Chung TP, Brownstein B, Stormo GD, Cobb JP: **Transcriptional profiles of human epithelial cells in response to heat: computational evidence for novel heat shock proteins.** *Shock (Augusta, Ga)* 2008, **29**(5):623-630.
 64. O'Brien ML, Cunningham ML, Spear BT, Glauert HP: **Peroxisome proliferators do not activate the transcription factors AP-1, early growth response-1, or heat shock factors 1 and 2 in rats or hamsters.** *Toxicol Sci* 2002, **69**(1):139-48.

doi:10.1186/1471-2164-11-16

Cite this article as: Vallanat et al: Analysis of the heat shock response in mouse liver reveals transcriptional dependence on the nuclear receptor peroxisome proliferator-activated receptor α (PPAR α). *BMC Genomics* 2010 **11**:16.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

