

RESEARCH ARTICLE

Open Access

Microarray Analysis of Response of *Salmonella* during Infection of HLA-B27- Transfected Human Macrophage-Like U937 Cells

Shichao Ge¹, Vittoria Danino², Qiushui He¹, Jay CD Hinton², Kaisa Granfors^{1*}

Abstract

Background: Human leukocyte antigen (HLA)-B27 is strongly associated with the development of reactive arthritis (ReA) in humans after salmonellosis. Human monocytic U937 cells transfected with HLA-B27 are less able to eliminate intracellular *Salmonella enterica* serovar Enteritidis than those transfected with control HLA antigens (e.g. HLA-A2). To investigate further the mechanisms by which HLA-B27-transfected cells allow increased replication of these bacteria, a DNA-based microarray was used for comparative genomic analysis of *S. Enteritidis* grown in HLA-B27- or HLA-A2-transfected cells. The microarray consisted of 5080 oligonucleotides from different serovars of *Salmonella* including *S. Enteritidis* PT4-specific genes. Bacterial RNA was isolated from the infected HLA-B27- or HLA-A2-transfected cells, reverse-transcribed to cDNA, and hybridized with the oligonucleotides on the microarrays. Some microarray results were confirmed by RT-PCR.

Results: When gene expression was compared between *Salmonella* grown in HLA-B27 cells and in HLA-A2 cells, 118 of the 4610 *S. Enteritidis*-related genes differed in expression at 8 h after infection, but no significant difference was detectable at 2 h after infection. These differentially expressed genes are mainly involved in *Salmonella* virulence, DNA replication, energy conversion and metabolism, and uptake and metabolism of nutrient substances, etc. The difference suggests HLA-B27-dependent modulation of *Salmonella* gene expression, resulting in increased *Salmonella* replication in HLA-B27-positive cells. Among the up-regulated genes were those located in *Salmonella* pathogenicity island (SPI)-2, which play a central role in intracellular survival and replication of *Salmonella*.

Conclusions: This is the first report to show the regulation of *Salmonella* gene expression by HLA-B27 during infection of host cells. This regulation probably leads to increased *Salmonella* survival and replication in HLA-B27-positive cells. SPI-2 genes seem to contribute significantly to the increased replication.

Background

The clinical outcomes of non-typhoidal salmonellosis range from self-limiting gastroenteritis to life-threatening systemic infections [1]. Many serovars of *Salmonella enterica* cause these infections, serovar Enteritidis being among the most common [2,3]. The acute gastrointestinal infection caused by *Salmonella* may result in complications such as reactive arthritis (ReA) [4-6]. Originally, ReA was described as an aseptic inflammation that develops after an infection elsewhere in the body [7]. ReA is an asymmetric polyarthritis and the

outcome of the disease ranges from mild symptoms to severe and chronic clinical manifestations. Up to 80% of patients with ReA express the HLA-B27 antigen [8,9].

Macrophages are important in the pathogenesis of *Salmonella* infections. They are an integral part of the immune response as they present antigens to the innate defence system and communicate with the adaptive immune system to resist the bacterial infection [10-12]. However, unlike many other pathogens, *Salmonella* can survive inside macrophages by adapting to this particular intracellular environmental niche. After *Salmonella* uptake into macrophages, the intracellular bacteria reside in large membrane-bound phagosomes, called spacious phagosomes (SP), which develop into *Salmonella*-containing vacuoles (SCV) [13]. The formation of

* Correspondence: kaisa.granfors@utu.fi

¹Department of Infectious Disease Surveillance and Control, National Institute for Health and Welfare, Turku, Finland
Full list of author information is available at the end of the article

SP or SCV favours the survival and replication of *Salmonella* in macrophages. Even so, *Salmonella* encounter intracellular host defence mechanisms, including reactive oxygen and nitrogen species (ROS and RNS), antimicrobial peptides, lysosomal enzymes, and adaptive immune responses [10,12]. In order to survive in the host and to avoid clearance by the host immune system, *Salmonella* express virulence factors to deal with this stressful environment [14].

Many virulence genes of pathogenic bacteria are located in large multigene chromosome regions termed pathogenicity islands (PAIs) [15]. In *Salmonella*, they are called *Salmonella* pathogenicity islands (SPIs) [16]. Two SPIs, SPI-1 and SPI-2, encode structurally similar but functionally distinct type III secretion systems (T3SS), specialized protein export machineries that *Salmonella* uses to deliver virulence proteins into the cytosol of host cells [17]. The SPI-1-encoded T3SS is active extracellularly. SPI-1 mediates invasion into non-phagocytic cells [18], and it is required for the intestinal inflammatory responses [19]. The SPI-2 virulence genes are expressed intracellularly and are required for the survival of bacteria in macrophages and systemic infections. SPI-2 mutant strains are dramatically attenuated, showing a 10^4 -fold reduction in virulence in LD₅₀ in the murine salmonellosis model [16] and impaired intracellular replication and survival in macrophages [20,21].

HLA-B27 confers a very strong genetic predisposition towards the development of a group of rheumatic disorders called spondyloarthropathies (SpA), including ankylosing spondylitis (AS) and ReA. HLA-B27-positive individuals have a five-fold higher incidence of ReA than the general population [22,23]. ReA occurs following certain infections, e.g. those caused by *Salmonella* and *Yersinia* pathogens [7,8,24,25]. Expression of HLA-B27 also increases the risk that the patient will suffer a more severe and prolonged disorder [6,8]. The interaction between ReA-triggering bacteria and HLA-B27-positive subjects is abnormal and leads to increased persistence of the causative microbes/microbial antigens in HLA-B27-positive patients [26-28]. The interaction between HLA-B27 molecules and arthritogenic microbes was investigated more thoroughly using *in vitro* infected cells. Experiments investigating the invasion of HLA-B27 cells by Gram-negative bacteria, including *Salmonella*, are inconclusive. Studies have shown either decreased [29] or similar [30] levels of invasion of HLA-B27-transfected murine L fibroblasts compared with control L cell lines (L cells transfected with other MHC class I genes) or increased invasion of HLA-B27-transfected intestinal epithelial Henle-407 cells by *Salmonella* [31]. Once inside the host cells, *Salmonella* is able to replicate more quickly [32,33], and is eliminated more slowly, in HLA-B27-positive cells

[30,34] than transfected control cells. The survival and persistence of *Salmonella* in the intracellular environment is associated with bacterial gene expression [35]. However, little is known about *Salmonella* gene expression in association with HLA-B27 during bacterial infection and persistence.

Results and Discussion

Cell surface expression of HLA-B27 and HLA-A2 molecules



The transfected HLA-B27 and HLA-A2 were always expressed on the surface of the respective cells, as detected by immunofluorescence in new batches of the cell lines (data not shown). The level of expression of the transfected molecules in U937 cells was similar to that of HLA-B51, one of the MHC class I molecules endogenously expressed by U937 cells [34]. In addition, the surface expression levels of the transfected molecules corresponded to the levels of those molecules endogenously expressed on peripheral blood monocytes [34].

Increased replication of *Salmonella* in HLA-B27-positive U937 cells

We used the *in vitro* model of infection established earlier in our laboratory to monitor growth of *S. Enteritidis* in macrophage-like U937 cells [32-34]. Cells transfected with HLA-B27 or HLA-A2 were infected with complement-opsonized salmonellae. HLA-A2 transfected cells were used as a negative control since it is a common tissue antigen that is not related to the development of ReA [34]. At 1, 4, 8, and 24 h post infection, the host cells were lysed and the number of living intracellular bacteria per cell was determined by counting the number of colony forming units (CFU) (Figure. 1). Consistent with our previous results, no difference in the uptake of *Salmonella* was observed between the two cell lines (1 h post infection) [34]. However, more bacteria were recovered from HLA-B27-expressing cells than HLA-A2-expressing cells at 8 h and even up to 24 h post infection, suggesting that HLA-B27-transfected cells become permissive for intracellular *Salmonella* survival and replication, which is also consistent with our previous studies [32-34].

Global gene expression profiles of *S. Enteritidis* in HLA-B27- and HLA-A2-transfected U937 cells


The 'SALSA' genomic *Salmonella* serovar microarray containing 5080 *Salmonella* oligonucleotides was used to monitor gene expression in *Salmonella* grown in the host cells [36]. Gene expression was compared between *Salmonella* grown in U937 cells and in LB broth. In total, 1,388 genes of *Salmonella* grown in HLA-B27 cells and 1,049 genes of *Salmonella* grown in HLA-A2 cells showed statistically significant differences in



expression at 2 h post infection when compared to the transcriptome in LB broth culture. At 8 h post infection, the expression of 1,352 genes of *Salmonella* grown in HLA-B27 cells and of 1,559 genes of *Salmonella* grown in HLA-A2 cells showed statistically significant differences from *Salmonella* grown in LB broth. The genes that showed significant expression differences and were up- or down-regulated more than two-fold for each individual comparison are listed in Additional file 1: Supplemental Tables 1-6. The categories of *Salmonella* genes showing differential expression between HLA-B27 cells and LB and between HLA-A2 cells and LB at 2 h post infection can be found in Figure. 2. In particular, the functional categories analysis showed large differences in the expression of genes required for cell motility and secretion, flagellae and chemotaxis and oxidative phosphorylation. These differences in gene expression reflect bacterial adaptation to the intracellular environments.

To confirm the microarray data, quantitative real time RT-PCR was performed on six genes selected from different functional categories and showing differential up-

or down-regulated expression between HLA-B27 and A2 cells: *mgfC*, *rpoE* and *ssaG* (up-regulated); *fliC*, *fumC* and *nuoA* (down-regulated) (Figure. 3). RT-PCR was performed using the same bacterial RNA from HLA-B27 cells, A2 cells and LB broth, as was used in microarray experiments. It is important that the genes selected covered the whole scale of the microarray data from high expression level to medium and to low expression level, as well as in differential expression from high fold changes to low fold changes in U937 cells vs. in LB, among up-regulated and down-regulated genes, respectively. The quantity of cDNA for each gene was obtained after normalization to the levels of *rfaH* cDNA, which was chosen as a control since *rfaH* is expressed stably at moderate levels under most of the conditions tested (see Additional file 2: Supplemental Table S8). The RT-PCR results of the six genes studied were consistent with the microarray data (Figure. 3), but due to the small number of genes studied with RT-PCR, data may be considered preliminary.



Salmonella gene expression is regulated by HLA-B27 at the 8 h time point

To determine whether bacterial gene expression plays a role in *Salmonella* replication/persistence in HLA-B27 cells, we assessed the transcriptome of *S. Enteritidis* in the two cell lines. Differential expression of the genes between HLA-B27 cells and HLA-A2 cells is based on the comparison of the gene expression in HLA-B27 cells to LB and the gene expression in HLA-A2 cells to LB. For example, for the gene whose expression is 6-fold in HLA-B27 cells compared to LB, and 2-fold in HLA-A2 cells compared to LB, the expression is 3-fold in HLA-B27 cells compared to HLA-A2 cells. No genes showed significant differences when the transcriptomes of *Salmonella* grown in HLA-B27- and HLA-A2-transfected cell lines were compared at 2 h post infection. This is consistent with the similarity observed in the intracellular

growth of *Salmonella* at this time point. In contrast, 118 genes showed significant differences in expression between *Salmonella* grown in HLA-B27 cells and in HLA-A2 cells at 8 h post infection (86 genes were up-regulated and 32 genes down-regulated (see Additional file 1: Supplemental Tables S6-1 and -2, Figure 4 and Table 1). Bacterial gene expression reflects the environmental conditions in which the bacteria reside. The difference between bacteria grown in HLA-B27 and HLA-A2 cells suggests that HLA-B27 modulates the intracellular environment and causes changes in *Salmonella* gene expression during macrophage infection, resulting in increased survival and replication in HLA-B27 cells. This modulation may be due to changes in host cell signalling pathways [33,37]. This is the first observation showing that *Salmonella* gene expression is influenced by HLA-B27. The analysis of functional categories revealed that the groups of genes most altered in the presence of HLA-B27 included those involved in *Salmonella* virulence, DNA replication, energy conversion and metabolism, and uptake and metabolism of nutrients.

Up-regulation of SPI-2 genes may prevent damage by ROS

SPI-2 genes have been shown to make *Salmonella* capable of growing intracellularly by avoiding killing by reactive oxygen species (ROS) [38]. During *Salmonella* replication in macrophages, the innate defence systems of the host cells are triggered to limit the infection. ROS produced by phagocyte NADPH oxidase (*phox*) are the most effective anti-bacterial agents [39]. NADPH oxidase catalyzes the univalent reduction of molecular oxygen and produces superoxide, which has modest




Table 1 The leading genes that were differentially expressed between the two cell lines


Name	Genbank	Function and product	Fold changes ^a
<i>ssaN</i>	AAL20339	Secretion system apparatus	3.97
<i>ssaO</i>	AAL20340	Secretion system apparatus	2.23
<i>ssaR</i>	AAL20343	Secretion system apparatus	4.09
<i>ssaS</i>	AAL20344	Secretion system apparatus	3.25
<i>ssaV</i>	AAL20338	Secretion system apparatus	2.56
<i>ssaA</i>	AAL20323	Secretion system apparatus	2.93
<i>sseI</i>	AAL19985	Gifsy-2 prophage; putative type III secreted protein	4.67
<i>sspH2</i>	AAL21143	Leucine-rich repeat protein, induced by the SPI-2 regulator <i>ssaR</i> /B	3.89
<i>dnaE</i>	AAL19195	DNA polymerase III, alpha subunit	2.68
<i>gyrA</i>	AAL21173	DNA gyrase, subunit A, type II topoisomerase	2.09
<i>gyrB</i>	AAL22694	DNA gyrase, subunit B, type II topoisomerase	2.39
<i>cyoC</i>	AAL19396	Cytochrome o ubiquinol oxidase subunit III	3.52
<i>cyoD</i>	AAL19395	Cytochrome o ubiquinol oxidase subunit IV	3.50
<i>nuoJ</i>	AAL21221	NADH dehydrogenase I chain J	2.19
<i>nuoK</i>	AAL21220	NADH dehydrogenase I chain K	2.42
<i>nuoM</i>	AAL21218	NADH dehydrogenase I chain M	2.39
<i>nuoN</i>	AAL21217	NADH dehydrogenase I chain N	2.27
<i>glnG</i>	AAL22844	Response regulator in two-component regulatory system with GlnL (EBP family)	2.88
<i>glnL</i>	AAL22845	Sensory kinase (phosphatase) in two-component regulatory system with GlnG	2.41
<i>glnA</i>	AAL22846	Glutamine synthetase	2.53
<i>gltB</i>	AAL22199	Glutamate synthase, large subunit	2.21
<i>gltD</i>	AAL22200	Glutamate synthase, small subunit	2.18
<i>aceF</i>	AAL19117	Pyruvate dehydrogenase, dihydrolipoyltransacetylase component	2.66
<i>fruA</i>	AAL21108	Sugar specific PTS system, fructose-specific transport	3.05
<i>fruF</i>	AAL21110	Phosphoenolpyruvate-dependent sugarphosphotransferase system, EIIA	2.67
<i>pykF</i>	AAL20302	Pyruvate kinase I (formerly F), fructose stimulated	3.33
<i>pstC</i>	AAL22714	ABC superfamily (membrane), high affinity phosphate	0.48

^a FDR (*P*-value < 0.05) was set up during microarray data analysis.

antibacterial activity but serves as a precursor for more toxic substrates, such as hydrogen peroxide and hydroxyl radicals [40]. Compared to those grown in HLA-A2 cells, *Salmonella* grown in HLA-B27 cells showed increased expression of SPI-2 genes including *ssaN*, *ssaO*, *ssaR*, *ssaS*, *ssaV*, *ssaA*, *sseI* and *sspH2* (Additional file 1: Supplemental Table S6-1). SPI-2 deficient mutants have impaired ability to grow in macrophages but are able to survive and are virulent within gp91*phox* knockout mice [39]. Patients with X-linked gp91*phox* mutations (chronic granulomatous disease, CGD) or gp91*phox* knockout mice that are unable to produce ROS are extremely susceptible to bacterial infection, including those caused by salmonellae [41,42]. Taken together, these findings indicated that SPI-2 is required for survival in the presence of ROS. The up-regulation of SPI-2 genes in HLA-B27 cells contributed to the increased intracellular growth and replication of *Salmonella*.

Intracellular growth of *Salmonella* SPI-2 mutants affected by HLA-B27

To verify the significance of SPI-2 genes in *Salmonella* replication in HLA-B27 cells, three genes most markedly up-regulated in HLA-B27 cells - *ssaN*, *ssaR* and *sspH2* - were deleted; *ssaN* and *ssaR* belong to the SPI-2 secretion apparatus and *sspH2* is an effector gene. The intracellular growth of the mutants was compared with that of wild-type *S. Enteritidis* PT4 KS8822/88. Mutation of *ssaR* dramatically impaired intracellular *Salmonella* growth in HLA-B27 cells, decreasing it 19-fold at the 8 h and 33-fold at the 24 h time points (Figure. 5A). It has previously been shown that Δ *ssaR* mutants cannot grow intracellularly in murine macrophages or form the elongated *Salmonella*-induced filaments (Sifs) in epithelial cells [43] required for the formation of *Salmonella*-containing vacuoles (SCV) [44]. The formation and maintenance of SCV promotes the survival and proliferation of bacteria in macrophages [45] and the lack of



formation of Sifs by *ssaR* mutants may account for the attenuation of this strain. The effector protein SspH2 is co-localized with vacuole-associated actin polymerization involved in the maintenance of SCV membrane integrity [46]. The *sspH2* mutant was also attenuated more than 4-fold in HLA-B27 cells at both 8 and 24 hours after infection (Figure. 5A). Unexpectedly, the

mutant carrying the *ssaN* deletion showed dramatically increased proliferation in HLA-B27 cells: 1.4-fold at the 8 h and 3.9-fold at the 24 h time points (Figure. 5A). The increased growth of the *ssaN* mutant in HLA-B27 cells might indicate that this mutant has good fitness in HLA-B27-positive U937 cells. The previous study by Eriksson et al [47] showed that the selection for *S. Typhimurium* mutants performed in murine macrophage-like J774-A.1 cells provided bacterial variants capable of a selective downregulation of nitric oxide (NO) expression. These variants showed increased host fitness. This is especially interesting, as in our model with murine L cells [30], iNOS activity and NO production were decreased in HLA-B27 positive cells, which correlated with increased *S. Enteritidis* growth. However, human U937 cells show very low iNOS activity and produce only small amounts of NO in general, and no difference between HLA-B27-positive and control cells was detected in this respect [48]. So the increased proliferation of the *ssaN* mutant in HLA-B27 cells, and no effect on growth in HLA-A2 cells (Figure 5B), perhaps indicates another specific and delicate relationship between *ssaN* and HLA-B27. These observations demonstrated a delicate balance between bacterial growth and pathogenesis.

Regulation of bacterial DNA synthesis by HLA-B27

As *Salmonella* replicates more in HLA-B27-transfected cells than in HLA-A2 transfectants, it is not surprising that genes involved in DNA synthesis are up-regulated in HLA-B27 cells compared to HLA-A2 cells. DNA polymerase III (pol III) holoenzyme is a major replicase responsible for DNA synthesis [49]. It is a multiprotein complex containing over 10 distinct subunits. The α subunit encoded by *dnaE* is one component of the catalytic core ($\alpha\epsilon\theta$) of the pol III holoenzyme and mutations affecting the α unit result in reduced growth of *S. enteritica* serovar Typhimurium [50]. The expression of the gene *dnaE* was induced 2.7-fold in HLA-B27 cells compared to HLA-A2 cells at 8 h time point (Table 1). Other *dna* genes including *dnaA* encoding DNA replication initial protein, *dnaB* encoding putative replicative DNA helicases, and *dnaN* encoding β -subunit of DNA polymerase III were up-regulated 1.9-, 1.4-, and 2.0-fold, respectively. However, *dnaQ* encoding ϵ -subunit of DNA polymerase III was expressed at the similar level in two cell lines (see Additional file 1: Supplemental Table S1). The up-regulation of most *dna* genes in HLA-B27 cells compared to HLA-A2 cells might indicate more DNA replication in HLA-B27 cells. Two genes (*gyrA* and *gyrB*) required for the production of the heterotetramer-DNA gyrase were also up-regulated in HLA-B27 cells compared with the control cells. DNA gyrase is an essential enzyme for bacterial DNA

synthesis; it introduces negative supercoils into DNA during replication [51]. The induction of *gyrA* and *gyrB* perhaps indicates that more DNA was synthesized by *Salmonella* resulting in increased bacterial proliferation in HLA-B27-transfected cells.

Salmonella substrate and energy metabolism

Growth in a nutrient-limited environment may be echoed in changes in gene expression. Expression of genes involved in substrate and energy metabolism, including *nuoJKMN* encoding NADH dehydrogenase I chains (J, K, M and N components), *cyoCD* encoding cytochrome O ubiquinol oxidase subunits III and IV, and *aceF* encoding a putative dehydrogenase was up-regulated in HLA-B27 cells compared to HLA-A2 cells. This suggests that more energy was produced in HLA-B27 cells under aerobic conditions, which could be connected to the enhanced bacterial replication observed in HLA-B27 cells [52,53].

Up-regulation of amino acid metabolism by *Salmonella* growth in HLA-B27 cells compared to HLA-A2 cells was evident from the induction of *gltB*, *gltD* and *glnA* encoding glutamate synthase/glutamine synthetase, and *glnGL* encoding a two-component regulator of glutamine synthetase [54]. Glutamate is an important precursor of other amino acids and of pyrimidine and purine synthesis in bacteria [55]. Moreover, sugar catabolism genes such as *aceF*, *fruA*, *fruF* and *pykF* were elevated in HLA-B27 cells. Collectively, these data suggest that HLA-B27 affects intracellular substrate metabolism and energy production by *Salmonella* during infection of U937 cells, probably due to the increased bacterial proliferation in these cells.

HLA-B27 modified ionic transport systems

Magnesium is involved in the stabilisation of the cellular membrane and functions as a coenzyme. MgtA and MgtB are two inducible transporters, and high up-regulation of *mgtB* and *mgtA* in both HLA-B27 and HLA-A2 macrophages might suggest that Mg^{2+} is extremely limited within the SCV of infected U937 cells [56,57]. From 1 to 8 h after infection, the degree to which these two genes were up-regulated was increased in both cell lines, indicating that the amounts of magnesium decreased during the course of infection. MgtC is also involved in Mg^{2+} transport in the intracellular survival of *Salmonella* and other bacteria in macrophages [58,59]. The two genes *mgtC* and *mgtB* are located in one operon, *mgtCB*, which belongs to SPI-3 on the *Salmonella* chromosome [58]. Our data showed that both *mgtC* and *mgtB* were less induced in HLA-B27-positive cells than in HLA-B27-negative cells, suggesting that HLA-B27 modulates the acquisition of Mg^{2+} in macrophages.

Phosphate availability is limited in macrophage vacuoles [60]. These observations are supported by our microarray data showing that the *phoBR* regulon, the *pstC* and *pstS* genes responsible for phosphate uptake and/or transport, were up-regulated during intracellular growth. The genes were induced to similar levels in both cell lines at 2 h after infection but the expression of *pstC* was down-regulated in HLA-B27 cells compared to HLA-A2 cells at 8 h postinfection. Taken together, these results suggest that HLA-B27 affects the intracellular ionic status of the SCV.

Conclusions

We have elucidated the global gene expression profile of *S. Enteritidis* PT4 KS8822/88 during intracellular growth in human monocyte/macrophages for the first time. The gene expression profile shown by *S. Enteritidis* during intracellular growth is very different from that exhibited in LB broth. However, it is similar to the patterns of gene expression reported for *S. Typhimurium* bacterial intracellular growth in macrophages [52]. Approximately one quarter of *S. Enteritidis* genes showed up- or down-regulation in host cells compared to LB broth. Among the genes up-regulated intracellularly were SPI-2 virulence genes, nutrient acquisition system genes, and ionic uptake and/or synthesis genes. Similar differences in intracellular *S. Typhimurium* gene expression were also seen between macrophages and medium *in vitro* [52]. SPI-2 genes are broadly induced to evade immune responses by host cells. Acquisition and transport system genes were elevated to take up the necessary elements. Among the down-regulated genes, some were related to central metabolism, such as the tricarboxylic acid cycle and oxidative phosphorylation, indicating that the level of oxygen is extremely low in the SCV intracellular environment, which seems contrary to the observation of intracellular *S. Typhimurium* growth [52]. Also, *Salmonella* metabolism and energy production were slower intracellularly than during growth in LB medium. These changes reflect the intracellular environment that the bacteria encounter during infection of macrophages [61].

The survival and proliferation of pathogens within macrophages is critical for establishing systemic infection, and this process is related to bacterial gene expression [62,63]. The expression of HLA-B27 in the U937 cells appeared to generate a favourable environment for intracellular *Salmonella* growth compared with HLA-A2-transfected cells, resulting in more bacteria in HLA-B27-transfected cells. The increased ability of *Salmonella* to survive is probably due to the altered gene expression. The expression of 118 genes at 8 h after infection differed significantly between HLA-B27 cells and HLA-A2 cells, suggesting that HLA-B27 modified *Salmonella* gene expression and/or affected the

intracellular bacterial growth environment during the infection of macrophages. Expression of SPI-2 genes, which play a key role in the increased persistence of *Salmonella* in macrophages, was up-regulated in HLA-B27 cells compared to HLA-A2 cells. DNA replication, energy production and nutrient metabolism were also increased, suggesting that more bacterial physiological activities occurred in HLA-B27 cells. This is consistent with the increased replication and survival of *Salmonella* in HLA-B27-transfected macrophages. Interestingly, expression of many SPI-2 genes, energy production and nutrient metabolism are also increased in epithelial cells where *Salmonella* replicates intracellularly [64], whereas e.g. Cyo and Cyd terminal oxidases, which use oxygen exclusively as terminal acceptor, are strongly downregulated in *Salmonella* inhabiting restrictive cells, fibroblasts [65] and HLA-A2 cells (Additional file 1: Supplemental Table S2-2 and S4-2).

Mutagenesis of the SPI-2 genes demonstrated the crucial role of SPI-2 in the survival and replication of *Salmonella* in HLA-B27 cells, as has been seen in other macrophages [18,19]. The intracellular replication and survival of the strains bearing deletion of either *ssaR* or *sspH2* was impaired in comparison with wild-type PT4 (Figure. 5A). Both proteins participate in the maintenance of the SCV membrane, which is important for the survival of *Salmonella* in infected macrophages. Increased growth of the *ssaN* mutant in HLA-B27-positive cells might indicate a special connection between this mutant and HLA-B27-positive cells, and may even be relevant to the pathogenesis of HLA-B27-associated ReA.

HLA-B27 modifies the host cell's signalling pathways during *Salmonella* infection. Our group recently demonstrated that mitogen activated protein kinase (MAPK) p38 is involved in *Salmonella* replication in U937 cells, and that the p38 MAPK pathway is deregulated in HLA-B27-transfected cells [33]. *Salmonella* uses type III secretion systems (T3SS) encoded in SPI-1 or -2 to deliver virulence proteins into host cells, which subsequently interfere with the host cell's signalling pathways [15,66,67]. Further experiments will be required to determine how *Salmonella* genes, particularly SPI-2 genes, affect intracellular signalling pathways and cause increased replication of bacteria in HLA-B27 cells. Nevertheless, these findings give new insights into the disturbed microbe-host interaction in HLA-B27-positive cells.

Methods

Cell lines and transfections

The human monocytic U937 cell line was obtained from ATCC (Rockville, USA). The full-length 6-kb genomic clone of human HLA-B*2705 DNA [68] in the vector

pUC19 and the full-length 5.1-kb genomic clone of human HLA-A2 DNA [69] in the vector pUC9 were kind gifts from Dr. Joel D. Taurog. For transfection, U937 cells were suspended in RPMI 1640 containing 1.8 mM L-glutamine and 1 mM sodium pyruvate (Gibco, Paisley, Scotland), and were then cotransfected by electroporation with the vectors carrying either HLA-B*2705 or HLA-A2 DNA with the plasmid pSV2neo (to provide resistance to geneticin [G-418]) (CalBiochem, Darmstadt, Germany). Stable transfectants were selected with 0.5 mg/ml geneticin and the expression of transfected HLA-B27 or -A2 molecules on the surface of U937 cells was examined by immunofluorescence (IF) and flow cytometry as described previously [34]. The transfected cell lines were maintained in RPMI 1640 supplemented with 10% heat-inactivated fetal calf serum (FCS; PAA laboratories, Linz, Austria), 1.8 mM L-glutamine and 50 µg/ml gentamicin (Gm) (both from Biological Institutes, Kibbutz Beit Herennek, Israel) at 37°C in a humidified atmosphere of 5% CO₂/95% air. Cell cultures were tested to confirm freedom from mycoplasma contamination.

Bacterial strains, plasmids and culture conditions

Bacterial strains and plasmids used in this study are listed in Table 2. The wild type *S. Enteritidis* strain was originally isolated from the stool of a patient with *Salmonella*-triggered ReA [34], and was typed as phage type 4 (PT4) and named *S. Enteritidis* PT4 KS8822/88. The *S. Enteritidis* and *Escherichia coli* strains were routinely grown in Luria-Bertani (LB) broth at 37°C. Bacteria carrying a temperature-sensitive plasmid (pKO-BEGA) were grown at 30°C. Media were supplemented with 100 µg/ml ampicillin (Amp) (Sigma) and 30 µg/ml kanamycin (Km) (Sigma) as required.

Cell infection model

For infection, the U937 cells were diluted to 1.0×10^6 cells/ml and then seeded in tissue culture flasks (75 cm²) or 24-well plates (Greiner, Germany). Prior to bacterial infection, the cells were cultured with phorbol myristate acetate (PMA; Sigma) for 24 h to differentiate them toward more mature macrophage-like cells. Two hours before infection, the adherent cells were washed with Hank's balanced salt solution (HBSS) and then overlaid with prewarmed RPMI 1640 supplemented with 10% human AB serum (Finnish Red Cross, Helsinki, Finland). The cells were then co-cultured with *Salmonella* at a multiplicity of infection (MOI) around 50:1. After 2 h of infection, the cells were washed three times with HBSS to remove non-adherent bacteria and overlaid with fresh RPMI 1640 containing 50 µg/ml Gm to kill extracellular bacteria. The *Salmonella*-infected cells were then incubated at 37°C for 1, 4, 8 or 24 hours as

Table 2 Bacterial strains and plasmids used in this study

Strains and plasmids	Characteristics	Source/reference
Strains		
S. Enteritidis PT4	Wild-type clinical isolate	34 ^a
PT4 (pKOBEGA)	PT4 containing helper plasmid pKOBEGA, Amp ^R	This study
PT4 Δ ssaN	PT4 Δ ssaN::Km, in-frame deletion, Km ^R	This study
PT4 Δ ssaR	PT4 Δ ssaR::Km, in-frame deletion, Km ^R	This study
PT4 Δ sspH2	PT4 Δ sspH2::Km, in-frame deletion, Km ^R	This study
E. coli MC4100	Used as template for amplification of kanamycin resistance gene	Gift from J. M. Ghigo
Plasmid pKOBEGA	Vector for recombination experiments, Amp ^R	Gift from J. M. Ghigo

^a : S. Enteritidis strain was typed as PT4 and named as the strain KS8822/88 in this study.

indicated. To determine the number of living intracellular bacteria, the infected cells were scraped using a cell scraper and the amount of living host cells was counted under a microscope after staining with Trypan blue. Host cells were lysed with 1% Triton X-100 in 1 × phosphate-buffered saline (PBS) at room temperature for 5-10 min, 20 µl of the lysate in ten-fold serial dilutions in PBS were added to LB agar plates, and the numbers of bacteria were reported as CFU (colony forming units).

RNA extraction

Two methods were used to extract bacterial RNA from infected cells. At 1 or 8 h post infection, the intracellular bacteria were recovered from infected macrophages and lysed with 1% Triton X-100 [34]. The lysate was first centrifuged at 250 × g at 4°C to remove pieces of broken host cells and then the supernatant was centrifuged at 12,000 × g for 5 min to precipitate bacterial cells. The approach developed for recovering bacteria from infected cells ensured the minimum contamination of extracted bacterial RNA with eukaryotic RNA. Otherwise, infected macrophages were lysed in 0.1% SDS, 1% acidic phenol, 19% ethanol in water on ice for 30 min [60] and similar results were obtained. Total bacterial RNA was extracted and purified using the Promega SV total RNA purification kit according to the manufacturer's instructions (Promega). Control RNAs from bacteria grown in LB broth to mid-logarithmic phase and from eukaryotic cells were isolated using the same RNA purification kit. RNA integrity was monitored by agarose gel electrophoresis and using the Agilent 2100 system (Agilent). The concentration and purity of RNA was measured by spectrophotometry Bio-RAD SmartSpec™3000 (Bio-Rad).

Microarray hybridisation

The 'SALSA' *Salmonella* serovar microarray was developed with 5080 *Salmonella* genes from different

serovars of *Salmonella* including 196 *S. Enteritidis* PT4 specific genes. More details are available at <http://www.ifr.bbsrc.ac.uk/safety/microarrays/#protocols>. A type II experiment design was used for hybridisation with labelled bacterial genomic DNA as a reference channel in each experiment [70]. The protocol for hybridisation was as previously described [52]. Briefly, total bacterial RNA (5 µg) from *Salmonella* grown in host cells or in LB broth was converted into cDNA using Superscript II reverse transcriptase and random primer as recommended by the manufacturer. The cDNA produced from the reverse transcription reaction and genomic DNA was subsequently fluorescently labelled using Cy5 or Cy3 dyes by random priming, with increasing labelling efficiency using Klenow enzyme. Labelled cDNA and genomic DNA were hybridized with the microarray slides overnight at 65°C. After hybridization, the slides were carefully washed, dried and scanned. All hybridizations were performed with at least three biological replicates.

Microarray data analysis

Fluorescent intensity data from each array were collected using a GenePix 4000A scanner (Axon Instruments). To compensate for unequal dye incorporation, data were centred by bringing the median natural logarithm of the ratios for each group of spots printed by the same pin to zero. The data were analyzed using GENE-SPRING™6 software (Silicon Genetics). The significance of the centred data at $P = 0.05$ was determined using a parametric-based statistical test, adjusting the individual P -value by the Benjamini and Hochberg false discovery rate multiple test correction.

Microarray data accession number

The supporting microarray data have been deposited in the Array Express database <http://www.ebi.ac.uk/arrayexpress> with accession number E-MEXP-1438.

Quantitative real time RT-PCR

Primer3 software <http://frodo.wi.mit.edu>[71] was used to design the primers for quantitative real time RT-PCR and purchased from Sigma (Table 3). The primers were designed to have similar melting temperatures and lengths of PCR products. About 1 µg of total RNA from the bacteria after 2 or 8 h infection of U937 cells or growth in LB medium was reverse-transcribed to 1st strand cDNA using AMV reverse transcriptase and the random primer provided in the kit following the manufacturer's instructions (1st Strand cDNA Synthesis Kit for RT-PCR (AMV)⁺, Roche). The resultant cDNA was used as a template during PCR, which was performed on a Roche LightCycler instrument and using the LightCycler FastStart DNA Master SYBR green I kit (Roche).

Table 3 Primers used in the real-time RT-PCR in this study

Genes	Primers	Sequences
<i>mgtC</i>	mgtC-F	5'-GTCTCTGGTATTGGCTTCTGG-3'
	mgtC-R	5'-TTGGCACAAAGAATAATGATCG-3'
<i>rpoE</i>	rpoE-K	5'-GACCGGATTGAAGCAGAAA-3'
	rpoE-R	5'-CCAGCTCCCGTAAGGTGAT-3'
<i>ssaG</i>	ssaG-F	5'-TTAGTGGATATGCTCTCCACA-3'
	ssaG-R	5'-TCATTTTATGATCAGTGAACITTCG-3'
<i>fliC</i>	fliC-F	5'-AGCCTGTCGCTGTTGACC-3'
	fliC-R	5'-CGCTGCAGTTGTGGTTG-3'
<i>fumC</i>	fumC-F	5'-CGGTATGGAACGCAAAGTG-3'
	fumC-R	5'-CTCCTGGCCTAAGGTGAGC-3'
<i>nuoA</i>	nuoA-F	5'-TGCTGCCTGATGCTGTA-3'
	unoA-R	5'-CGCTTTCGCGGATAGAAG-3'
<i>rfaH</i>	rfaH-F	5'-ACTTCAGCGTGCTCAGGAA-3'
	rfaH-R	5'-GCGTGCGCTGATTGTAGT-3'

The protocol for the PCR reaction was an initial denaturation at 95°C for 10 min, followed by 40 cycles of denaturation at 95°C for 5 s, annealing at 56-64°C for 5 s and extension at 72°C for 10 s. For certain pairs of primers the annealing temperatures might be changed to optimize the yield of products. The amplified specific products were examined by melting temperature curves and agarose gel electrophoresis. Relative gene expression was quantified using a standard curve method plotted from ten-fold serial dilutions of known quantities of cDNA samples. Each standard curve for each gene was then used to transform threshold cycle (Ct) to the relative amount of unknown cDNA of the gene. The analysis was based on the Ct values of the specific genes, with *rfaH* as a normalizer. The amount of the specific gene was divided by the amount of *rfaH* in the same cDNA preparation to calculate the normalized value. Relative gene expression was presented as ratios of the normalized values of cDNA from bacterial growth in U937 cells to the growth in LB broth. All experiments were performed in triplicate using independent RNA preparations (biological replicates).

DNA manipulations and preparation of electrocompetent cells

Plasmid DNA from *E. coli* was purified with a Quantum Prep plasmid miniprep kit (Bio-Rad). Plasmids were transformed into the strain *S. Enteritidis* PT4 KS8822/88 by electroporation. Transformants carrying the Red helper plasmid were made electrocompetent as described previously [72] with some modifications. Cells were grown overnight in LB broth Amp at 30°C and then 1 ml of overnight cultures were used to inoculate 100 ml of LB broth Amp and incubation was continued to an OD₆₀₀ of 0.15-0.2. L-arabinose (Sigma) was then

added to a final concentration of 10 mM and incubation was continued until the OD₆₀₀ reached 0.7. The suspension was cooled on ice for 20 min, and the cells were made electrocompetent by washing twice with the same volume of ice-cold water and then once with 40 ml of ice-cold 10% glycerol. The cells were finally resuspended in 1 ml of ice-cold 10% glycerol, then divided into 50 µl working aliquots and kept at -70°C for up to two months. Total genomic DNA was isolated according to the manufacturer's instructions (High Pure PCR template Preparation kit, Roche).

Construction of gene mutants by the λ Red mutagenesis method

Salmonella genes were disrupted by the method described previously [72,73]. Briefly, purified plasmid pKOBEGA was introduced into *Salmonella* by electroporation, and transformants were selected on LB agar Amp after incubation for 24 h at 30°C. The λ Red

Table 4 Primers used in the gene mutagenesis in this study

Genes	Primers	Sequences	
<i>ssaN</i>	ssaN-Km-F	5'-CTTTGCTATCTCCTTTTACGAGTACAATCGGGCTTCACTGC GGGCAGCAAGTGATGGCCTAAAGCCACGTTGTGTCTCAA- 3' ^α	
	ssaN-Km-R	5'-CTTCAGACAGTGATAAATCGATGAATTCGCGGACTTCTCGT CCACGTTACCAATTAACAGCGCTGAGGTTCTGCCTCGTG- 3' ^α	
	ssaN-F	5'-CGTTGTTAAATGCGTGGTTG-3'	
	ssaN-R	5'-CCCATTCCCGTACGTTCTAA-3'	
	<i>ssaR</i>	ssaR-Km-F	5'-GCGCTTGACTTTCCTTATTCAATTATGGGGCCGACGCTATT AGCTGTAAAAGAGCGCTGAAAGCCACGTTGTGTCTCAA- 3' ^α
		ssaR-Km-R	5'-ATTAATATGAGCAAAGAATCAGGTTTTATCTTTCTTTTAT GTTCTTCAGGCCAGGTTCCGGCCTGAGGTTCTGCCTCGTG- 3' ^α
	ssaR-F	5'-TATCGCACTGTATGGCCTTG-3'	
	ssaR-R	5'-ACCGCTGCCAGTAAAATA-3'	
<i>sspH2</i>	sspH2-Km-F	5'-CTTTATGAAGTTTTCCGTCTCACTCAGTCTGTCCAGGAAGA GGCTGAATGCGTCGGCGTTAAAGCCACGTTGTGTCTCAA- 3' ^α	
	sspH2-Km-R	5'-ACACTGGTTATTCTGATAATAATCTGACCAGCCTGCCGGC GCTGCCGCCAGAAGCTGCGGCGCTGAGGTTGTGTCTCAA- 3' ^α	
	sspH2-F	5'-TCATCTTCAGCCAGTTGTGC-3'	
	sspH2-R	5'-GTAATCGCCGATTTATCGT-3'	

^α: 80-nucleotide (nt)-long primers including 60 nt homology extensions complementary to the targeted regions of the genes and 20 nt priming sequences (underlined) for the synthesis of the kanamycin resistance cassette gene from *E. coli* MC4100 *ybeW::Km*.

helper plasmid pKOBEGA is a low-copy-number plasmid that contains an ampicillin resistance gene, a temperature-sensitive origin of replication and the Red system, including three genes expressing the Exo, Bet and Gam functions of phage λ , which helps allelic exchanges between linear DNA and the corresponding region on the chromosome [74]. The *Salmonella* strains carrying the λ Red helper plasmid were made electrocompetent as described above. PCR linear DNA fragments were generated with specific hybrid primers and high-fidelity thermophilic DNA polymerase (Dynazyme Ext; Finnzymes). The primer pairs shown in Table 4 were used. PCR products were purified with a QIAquick PCR Purification kit (QIAGEN) and 5 μ g of purified product (5-10 μ l) was electroporated (25 μ F, 200 Ω , 2.5 kV) into 50 μ l of electrocompetent cells according to the manufacturer's instructions (Bio-Rad), then 950 μ l of LB broth was added to the shocked cells and the solution was transferred into clear eppendorf tubes and incubated for 2 h at 37°C, and then spread on to LB agar Km to select Km transformants after overnight growth at 37°C. Selected antibiotic resistance colonies were then grown on LB broth Km at 43°C for 24 h and then spread on LB agar Km or Amp at 37°C to check the loss of the helper plasmid. The mutated genes were confirmed by PCR using the primers shown in Table 4, which are external to the site of mutagenesis with the internal deletion replaced by the kanamycin cassette.

Additional material

Additional file 1: Supplemental Tables S1- 6 - Global gene expression of *S. Enteritidis* PT4 in HLA-B27- or HLA-A2-transfected cells using microarray assay. The Supplemental Tables S1-6 contain gene expression profiles of *S. Enteritidis* PT4 strain during infection of U937 cells (Table S1), genes up- or down-regulated more than two-fold in HLA-A2-transfected U937 cells compared to LB culture at 2 h after infection (Tables S2-1, -2), in HLA-B27-transfected cells compared to LB culture at 2 h after infection (Table S3-1,-2), in HLA-A2-transfected U937 cells compared to LB culture at 8 h after infection (Tables S4-1, -2), in HLA-B27-transfected cells compared to LB culture at 8 h after infection (Table S5-1,-2) and in HLA-B27-transfected cells compared to HLA-A2-transfected cells at 8 h after infection (Tables S6-1, -2).

Additional file 2: Supplemental Tables S7 and S8 - The correlation analysis between PT-PCR and array data and the expression of the gene *rfaH* in different conditions. The Supplemental Tables S7 and S8 contain the correlation analysis between PT-PCR and array data (Table S7) and the expression of the gene *rfaH* in different conditions (Table S8).

Acknowledgements

We are grateful for useful discussions and advice from Jari Jalava, Yangyang Wang, Sanna Vähämäki, Anna Sahlberg, Isabelle Hautefort and Arthur Thompson. We thank Tuija Turjas, Tuula Rantasalo, Päivi Haaranen, and Erkki Nieminen for their technical assistance. We also thank Anja Siitonen for typing our *S. Enteritidis* strain, Joel D. Taurig for providing the HLA-B27 and HLA-A2 genomic DNA, and Jean-Marc Ghigo and Iñigo Lasa for providing *E.*

coli MC4100 *ybeW::Km*, plasmid pKOBEGA and bacteriophage P22. Biomedes is acknowledged for copyediting of the manuscript.

This study was supported by grants from the Academy of Finland and the Sigrid Jusélius Foundation to Kaisa Granfors and the BBSRC core strategic grant to Jay C. D. Hinton and the DEFRA VTRI VT104 grant to Vittoria Danino.

Author details

¹Department of Infectious Disease Surveillance and Control, National Institute for Health and Welfare, Turku, Finland. ²Molecular Microbiology Group, Institute of Food Research, Colney, Norwich, UK.

Authors' contributions

SCG designed and carried out the experiments, and wrote the manuscript. VD performed the microarray tests and analyzed the data. QSH designed the experiments and analyzed the data. JCDH conceived and designed the experiments. KG conceived and designed the experiments, and wrote the manuscript. All authors reviewed the manuscript.

Received: 3 December 2009 Accepted: 30 July 2010

Published: 30 July 2010

References

1. Fierer J, Guiney DG: **Diverse virulence traits underlying different clinical outcomes of *Salmonella* infection.** *J Clin Invest* 2001, **107**:775-780.
2. Mishu B, Koehler J, Lee LA, Rodrigue DF, Brenner H, et al: **Outbreaks of *S. Enteritidis* infections in the United States, 1985-1991.** *J Infect Dis* 1994, **169**:547-552.
3. Rabsch W, Tschäpe H, Bäuml AJ: **Non-typhoidal salmonellosis: emerging problems.** *Microbes and Infect* 2001, **3**:237-247.
4. Burmester GR, Daser A, Kamradt T, Krause A, Mitchison NA, et al: **Immunology of reactive arthritis.** *Annu Rev Immunol* 1995, **13**:229-250.
5. Locht H, Molbak K, Krogfelt KA: **High frequency of reactive joint symptoms after an outbreak of *Salmonella* Enteritidis.** *J Rheumatol* 2002, **29**:767-77.
6. Ekman P, Kiveskari J, Granfors K: **Modification of disease outcome in *Salmonella*-infected patients by HLA-B27.** *Arthritis Rheum* 2000, **43**:1527-34.
7. Ahvonen P, Sievers K, Aho K: **Arthritis associated with *Yersinia enterocolitica* infection.** *Acta Rheumatol Scand* 1969, **15**:232-253.
8. Leirisalo M, Skylv G, Kousa M, Voipio-Pulkki L, Suoranta H, et al: **Followup study on patients with Reiter's disease and reactive arthritis, with special reference to HLA-B27.** *Arthritis Rheum* 1982, **25**:249-259.
9. Aho K, Ahvonen P, Lassus A, Sievers K, Yliliskainen A: **HLA-A in reactive arthritis: a study of *Yersinia* arthritis and Reiter's disease.** *Arthritis Rheum* 1974, **17**:521-526.
10. Mastroeni P: **Immunity to systemic *Salmonella* infections.** *Curr Mol Med* 2002, **2**:393-406.
11. Nicolas P, Mor A: **Peptides as weapons against microorganisms in the chemical defense system of vertebrates.** *Annu Rev Microbiol* 1995, **49**:277-304.
12. Vazquez-Torres A, Jones-Carson J, Mastroeni P, Ischiropoulos H, Fang FC: **Antimicrobial actions of the NADPH phagocyte oxidase and inducible nitric oxide synthase in experimental salmonellosis. I. Effects on microbial killing by activated peritoneal macrophages *in vitro*.** *J Exp Med* 2000, **192**:227-236.
13. Alpuche-Aranda CM, Racoosin EL, Swanson JA, Miller SI: ***Salmonella* stimulate macrophage macropinocytosis and persist within spacious phagosomes.** *J Exp Med* 1994, **179**:601-608.
14. Ernst RK, Guina T, Miller SI: **How intracellular bacteria survive: surface modification that promotes resistance to host innate immune responses.** *J Infect Dis* 1999, **179**:S326-330.
15. Hacker J, Blum-Oehler G, Muhldorfer I, Tschäpe H: **Pathogenicity island of virulent bacteria: structure, function and impact on microbial evolution.** *Mol Microbiol* 1997, **23**:1089-1097.
16. Shea JE, Hensel M, Gleeson C, Holden DW: **Identification of a virulence locus encoding a second type III secretion system in *Salmonella typhimurium*.** *Proc Natl Acad Sci USA* 1996, **93**:2593-2597.
17. Hueck CJ: **Type III protein secretion systems in bacterial pathogens of animals and plants.** *Microbiol. Mol Biol Rev* 1998, **62**:379-433.

18. Galán JE: Molecular genetic bases of *Salmonella* entry into host cells. *Mol Microbiol* 1996, 20:263-271.
19. Wallis TS, Galyov EE: Molecular basis of *Salmonella*-induced enteritis. *Mol Microbiol* 2000, 36:997-1005.
20. Cirillo DM, Valdivia RH, Monack DM, Falkow S: Macrophage-dependent induction of the *Salmonella* pathogenicity island 2 type III secretion system and its role in intracellular survival. *Mol Microbiol* 1998, 30:175-188.
21. Hensel M, Shea JE, Waterman SR, Mundy R, Nikolaus T, Banks G, Gleeson C, Fang FC, Holden DW: Genes encoding putative effector proteins of the type III secretion systems of *Salmonella* pathogenicity islands 2 are required for bacterial virulence and proliferation in macrophages. *Mol Microbiol* 1998, 30:163-174.
22. Colmegna I, Cuchacovich R, Espinoza LR: HLA-B27-associated reactive arthritis: pathogenetic and clinical considerations. *Clin Microbiol Rev* 2004, 17:348-369.
23. Feltkamp TEW: Factors involved in the pathogenesis of HLA-B27 associated arthritis. *Scand J Rheumatol* 1995, 24(Suppl 101):213-217.
24. Rohekar S, Tsui FW, Tsui HW, Xi N, Riarh R, Bilotta R, Inman RD: Symptomatic acute reactive arthritis after an outbreak of salmonella. *J Rheumatol* 2008, 35:1599-1602.
25. Mäki-Ikola O, Granfors K: *Salmonella*-triggered reactive arthritis. *Lancet* 1992, 339:1096-1098.
26. Granfors K, Merilähti-Palo R, Luukkainen R, Möttönen T, Lahesmaa R, et al: Persistence of *Yersinia* antigens in peripheral blood cells from patients with *Yersinia enterocolitica* O:3 infection with or without reactive arthritis. *Arthritis Rheum* 1998, 41:855-862.
27. Granfors K, Jalkanen S, Mäki-Ikola O, Lahesmaa-Rantala R, Saario R, et al: *Salmonella* lipopolysaccharide in synovial cells from patients with reactive arthritis. *Lancet* 1990, 335:685-688.
28. Granfors K, Jalkanen S, von Essen R, Lahesmaa-Rantala R, Isomaki O, et al: *Yersinia* antigens in synovial-fluid from patients with reactive arthritis. *N Engl J Med* 1989, 320:216-221.
29. Kapasi K, Inman RD: HLA-B27 expression modulates gram-negative bacterial invasion into transfected L cells. *J Immunol* 1992, 148:3554-3559.
30. Virtala M, Kirveskari J, Granfors K: HLA-B27 modulates the survival of *Salmonella enteritidis* in transfected L cells, possibly by impaired nitric oxide production. *Infect Immun* 1997, 65:4236-4242.
31. Saarinen M, Ekman P, Ikeda M, Virtala M, Grönberg A, et al: Invasion of *Salmonella* into human intestinal epithelial cells is modulated by HLA-B27. *Rheumatology* 2002, 41:651-657.
32. Penttinen MA, Heiskanen KM, Mohapatra R, DeLay ML, Colbert RA, et al: Enhanced intracellular replication of *Salmonella* Enteritidis in HLA-B27-expressing human monocytic cells: dependency on glutamic acid 45 in the B pocket of HLA-B27. *Arthritis Rheum* 2004, 50:2255-2263.
33. Sahlberg A, Penttinen MA, Keiskanen KM, Colbert RA, Sistonen L, et al: Evidence that the p38 MAP kinase pathway is dysregulated in HLA-B27-expressing human monocytic cells: correlation with HLA-B27 misfolding. *Arthritis Rheum* 2007, 56:2652-2662.
34. Laitio P, Virtala M, Salmi M, Pelliniemi LJ, Yu DT, et al: HLA-B27 modulates intracellular survival of *Salmonella enteritidis* in human monocytic cells. *Eur J Immunol* 1997, 27:1331-1338.
35. Buchmeier NA, Heffron F: Induction of *Salmonella* stress proteins upon infection of macrophages. *Science* 1990, 248:730-732.
36. Karavolos MH, Bulmer DM, Thompson A, Winzer K, Williams P, Hinton JC, Khan CM: Adrenaline modulates the global transcriptional profiles of *Salmonella* revealing a role in the antimicrobial peptide and oxidative stress resistance responses. *BMC Genomics* 2008, 9:458-471.
37. Ikawa T, Ikeda M, Yamaguchi A, Tsai WC, Tamura N, et al: Expression of arthritis-causing HLA-B27 on HeLa cells promotes induction of c-fos in response to *in vitro* invasion by *Salmonella typhimurium*. *J Clin Invest* 1998, 101:263-272.
38. Vazquez-Torres A, Xu Y, Jones-Carson J, Holden DW, Lucia SM, et al: *Salmonella* pathogenicity island 2-dependent evasion of the phagocyte NADPH oxidase. *Science* 2000, 287:1655-1658.
39. Vazquez-Torres A, Fang FC: *Salmonella* evasion of the NADPH phagocyte oxidase. *Microbes Infect* 2001, 3:1313-1320.
40. Babior BM: The respiratory burst oxidase. *Curr Opin Hematol* 1995, 2:55-60.
41. Mouy R, Fischer A, Vilmer E, Seger R, Griscelli C: Incidence, severity, and prevention of infections in chronic granulomatous disease. *J Pediatr* 1989, 114:555-560.
42. Shiloh MU, MacMicking JD, Nicholson S, Brause JE, Potter S, et al: Phenotype of mice and macrophages deficient in both phagocyte oxidase and inducible nitric oxide synthase. *Immunity* 1999, 10:29-38.
43. Brumell JH, Rosenberger CM, Gotto GT, Marcus SL, Finlay BB: SifA permits survival and replication of *Salmonella typhimurium* in murine macrophages. *Cell Microbiol* 2001, 3:75-84.
44. Garcia-del Portillo F, Zwick MB, Leung KY, Finlay BB: *Salmonella* induces the formation of filamentous structures containing lysosomal membrane glycoproteins in epithelial cells. *Proc Natl Acad Sci USA* 1993, 90:10544-10548.
45. Knodler LA, Steele-Mortimer O: Taking possession: biogenesis of the *Salmonella*-containing vacuole. *Traffic* 2003, 4:587-599.
46. Méresse S, Unsworth KE, Habermann A, Griffiths G, Fang FC, et al: Remodelling of the actin cytoskeleton is essential for replication of intravacuolar *Salmonella*. *Cell Microbiol* 2001, 3:567-577.
47. Eriksson S, Björkman J, Borg S, Syk A, Pettersson S, Andersson DI, Rhen M: *Salmonella typhimurium* mutants that downregulate phagocyte nitric oxide production. *Cell Microbiol* 2000, 2:239-250.
48. Ekman P, Saarinen M, He Q, Virtala M, Salmi M, Granfors K: Human monocytic U937 cells kill *Salmonella* in vitro by NO-independent mechanism. *Infect Immun* 1999, 67:3670-3673.
49. Burgers PM, Kornberg A, Sakakibara Y: The *dnaN* gene codes for the β subunit of DNA polymerase III holoenzyme of *Escherichia coli*. *Proc Natl Acad Sci USA* 1981, 78:5391-5395.
50. Lancy ED, Lifshits MR, Munson P, Maurer R: Nucleotide sequences of *dnaE*, the gene for the polymerase subunit of DNA polymerase III in *Salmonella typhimurium*, and a variant that facilitates growth in the absence of another polymerase subunit. *J Bacteriol* 1989, 171:5581-5586.
51. Rudd KE, Menzel R: *his* operons of *Escherichia coli* and *Salmonella typhimurium* are regulated by DNA supercoiling. *Proc Natl Acad Sci USA* 1987, 84:517-521.
52. Eriksson S, Lucchini S, Thompson A, Rhen M, Hinton JCD: Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*. *Mol Microbiol* 2003, 47:103-118.
53. Garcia-del Portillo F: *Salmonella* intracellular proliferation: where, when and how? *Microbes Infect* 3:1305-1311.
54. Reitzer LJ, Magasanil B: Asparagine synthetases of *Klebsiella aerogenes*: properties and regulation of synthesis. *J Bacteriol* 1982, 151:1299-1313.
55. Boyce JD, Wilkie I, Harper M, Paustian ML, Kapur V, et al: Genomic scale analysis of *Pasteurella multocida* gene expression during growth within the natural chicken host. *Infect Immun* 2002, 70:6871-6879.
56. Smith RL, Kaczmarek MT, Kucharski LM, Maguire ME: Magnesium transport in *Salmonella typhimurium*: regulation of *mgtA* and *mgtCB* during invasion of epithelial and macrophage cells. *Microbiology* 1998, 144:1835-1843.
57. Snavely MD, Miller CG, Maguire ME: The *mgtB* Mg²⁺ transport locus of *Salmonella typhimurium* encodes a P-type ATPase. *J Biol Chem* 1991, 266:815-823.
58. Blanc-Potard AB, Groisman EA: The *Salmonella selC* locus contains a pathogenicity island mediating intramacrophage survival. *EMBO J* 1997, 16:5376-5385.
59. Maloney KE, Valvano MA: The *mgtC* gene of *Burkholderia cenocepacia* is required for growth under magnesium limitation conditions and intracellular survival in macrophages. *Infect Immun* 2006, 74:5477-5486.
60. Deiwick J, Nikolaus T, Erdogan S, Hensel M: Environmental regulation of *Salmonella* pathogenicity island 2 gene expression. *Mol Microbiol* 1999, 31:1759-1773.
61. Buchmeier NA, Heffron F: Induction of *Salmonella* stress proteins upon infection of macrophages. *Science* 1990, 248:730-732.
62. Fields PI, Swanson RV, Haidaris CG, Heffron F: Mutants of *Salmonella typhimurium* that cannot survive within the macrophage are avirulent. *Proc Natl Acad Sci USA* 1986, 83:5189-5193.
63. Ohl ME, Miller SI: *Salmonella*: a model for bacterial pathogenesis. *Annu Rev Med* 2001, 52:259-274.
64. Hautefort I, Thompson A, Eriksson-Ygberg S, Parker ML, Lucchini S, Danino V, Bongaerts RJ, Ahmad N, Rhen M, Hinton JC: During infection of epithelial cells *Salmonella enterica* serovar Typhimurium undergoes a time-dependent transcriptional adaptation that results in simultaneous expression of three type 3 secretion systems. *Cell Microbiol* 2008, 10:958-984.

65. García-del Portillo F, Núñez-Hernández C, Eisman B, Ramos-Vivas J: **Growth control in the *Salmonella* -containing vacuole.** *Curr Opin Microbiol* 2008, **11**:46-52.
66. Meccas J, Strauss EJ: **Molecular mechanisms of bacterial virulence: Type III secretion and pathogenicity islands.** *Emerg Infect Dis* 1996, **30**:271-288.
67. Uchiya K, Groisman EA, Nikai T: **Involvement of *Salmonella* pathogenicity island 2 in the up-regulation of interleukin-10 expression in macrophages: role of protein kinase A signal pathway.** *Infect Immun* 2004, **72**:1964-1973.
68. Taurog JD, Lowen L, Forman J, Hammer RE: **HLA-B27 in inbred and non-inbred transgenic mice: cell surface expression and recognition as an alloantigen in the absence of human β_2 -microglobulin.** *J Immunol* 1988, **141**:4020-4023.
69. Koller BH, Orr HT: **Cloning and complete sequence of an HLA-A2 gene: analysis of two HLA-A alleles at the nucleotide level.** *Immunol* 1985, **134**:727-733.
70. Yang YH, Speed T: **Design issues for cDNA microarray experiments.** *Nat Rev Genet* 2002, **3**:579-588.
71. Rozen S, Skaletsky H: **Primer3 on the WWW for general users and for biological programmers.** *Methods Mol Biol* 2000, **132**:365-386.
72. Latasa C, Roux A, Toledo-Arana A, Ghigo JM, Gamazo C, et al: **BapA, a large secreted protein required for biofilm formation and host colonization of *Salmonella enterica* serovar Enteritidis.** *Mol Microbiol* 2005, **58**:1322-1339.
73. Datsenko KA, Wanner BL: **One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products.** *Proc Natl Acad Sci USA* 2000, **97**:6640-6645.
74. Chaverche MK, Ghigo JM, d'Enfert C: **A rapid method for efficient gene replacement in the filamentous fungus *Aspergillus nidulans*.** *Nucleic Acids Res* 2000, **28**:E97.

doi:10.1186/1471-2164-11-456

Cite this article as: Ge et al.: Microarray Analysis of Response of *Salmonella* during Infection of HLA-B27- Transfected Human Macrophage-Like U937 Cells. *BMC Genomics* 2010 **11**:456.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

